



Document Title

**Summary of the fate and behaviour in the environment  
fluoxastrobin + prothioconazole EC 200 (100+100 g/L)**

Data Requirements

**EU Regulation 1107/2009 & EU Regulation 284/2013**

**Document MCP**

**Section 9: Fate and behaviour in the environment**

According to the guidance document, SANCO/10181/2013, for preparing dossiers for the approval of a chemical active substance

Date

**2016-04-12**

Author(s)

[Redacted]

**Bayer CropScience**



M-542831-01-2

*This document is copyright protected.  
Any distribution, reproduction, publication requires  
the consent of Bayer AG (or its respective affiliate or  
any other commercial purpose is prohibited and constitutes  
a violation of the underlying license agreement.*



## OWNERSHIP STATEMENT

This document, the data contained in it and copyright therein are owned by Bayer CropScience. No part of the document or any information contained therein may be disclosed to any third party without the prior written authorisation of Bayer CropScience.

The summaries and evaluations contained in this document are based on unpublished proprietary data submitted for the purpose of the assessment undertaken by the regulatory authority. Other registration authorities should not grant, amend, or renew a registration on the basis of the summaries and evaluation of unpublished proprietary data contained in this document unless they have received the data on which the summaries and evaluation are based, either:

- From Bayer CropScience; or
- From other applicants once the period of data protection has expired.

This document is copyright protected and its content for regulatory purposes is confidential. Any distribution, reproduction or publication of this document without the prior written consent of Bayer AG (or its respective regulatory authorities) is prohibited and constitutes a violation of the underlying license agreement.



### Version history

Date	Data points containing amendments or additions <sup>1</sup> and brief description	Document identifier and version number

<sup>1</sup> It is suggested that applicants adopt a similar approach to showing revisions and version history as outlined in SANCO/10180/2013 Chapter 4 How to revise an Assessment Report

*This document is copyright protected (requires the consent of Bayer AG (or its respective affiliate) or any other commercial purpose is prohibited and constitutes a violation of the underlying license agreement.*



Table of Contents

	Page
CP 9	FATE AND BEHAVIOUR IN THE ENVIRONMENT..... 5
CP 9.1	Fate and behaviour in soil..... 8
CP 9.1.1	Rate of degradation in soil..... 8
CP 9.1.1.1	Laboratory studies ..... 8
CP 9.1.1.2	Field studies ..... 8
CP 9.1.1.2.1	Soil dissipation studies ..... 9
CP 9.1.1.2.2	Soil accumulation studies ..... 9
CP 9.1.2	Mobility in the soil ..... 9
CP 9.1.2.1	Laboratory studies ..... 9
CP 9.1.2.2	Lysimeter studies ..... 9
CP 9.1.2.3	Field leaching studies ..... 9
CP 9.1.3	Estimation of concentrations in soil ..... 10
CP 9.2	Fate and behaviour in water and sediment ..... 16
CP 9.2.1	Aerobic mineralisation in surface water ..... 17
CP 9.2.2	Water/sediment study ..... 17
CP 9.2.3	Irradiated water/sediment study ..... 17
CP 9.2.4	Estimation of concentrations in groundwater ..... 18
CP 9.2.4.1	Calculation of concentrations in groundwater ..... 19
CP 9.2.4.2	Additional field tests ..... 31
CP 9.2.5	Estimation of concentrations in surface water and sediment ..... 32
CP 9.3	Fate and behaviour in air ..... 53
CP 9.3.1	Route and rate of degradation in air and transport via air ..... 53
CP 9.4	Estimation of concentrations for other routes of exposure ..... 53

This document is copyright protected. Any distribution, reproduction or publication requires the consent of Bayer AG (or its respective affiliate). Any use of the document for regulatory or constitutive purposes is prohibited and constitutes a violation of the underlying license agreement.



CP 9 FATE AND BEHAVIOUR IN THE ENVIRONMENT

Use patterns considered in this risk assessment

Table CP 9- 1: Intended application patterns

Crop	Timing of application (range)	Number of applications	Application interval [days]	Maximum label rate per treatment [kg/ha]	Application rate per treatment [g/ha]	
					Fluoxastrobin	Prothioconazole
Wheat, rye, triticale*	BBCH 30-69	1-2	14-21	1.5	150	150
Barley, oats*	BBCH 30-61	1-2	14-21	1	125	125
Onions**	BBCH 15-47	1-2	10	1.0-1.25	100-125	100-125

\* Use in Central Europe; \*\* Use in Southern Europe






Compounds addressed in this document

In addition to the active substance fluoxastrobin, the degradation products summarised in Table CP 9- 2 are addressed in this document as they were major in environmental fate studies. In this paragraph the approach to the risk assessment of the Z-isomer of fluoxastrobin is specifically considered. The chemical structure of fluoxastrobin contains an oxime ether moiety. Due to the substitution pattern of that double bond E- and Z-isomers exist. The common name fluoxastrobin denotes the E-isomer. The Z-isomer is known to be an impurity in technical fluoxastrobin (specification limit 2 mg/kg). The Z-isomer can be formed from the E-isomer by photolytic processes exclusively. The transformation will lead to an equilibrium state in which the E-isomer is the more stable and energetically preferred isomer (ratio in aqueous solution about 10:1 = E / Z). In the environment the Z-isomer shows very similar degradation behaviour and a better soil sorption than the E-isomer. Further, the Z-isomer shows a very similar toxicological profile. A study with *Daphnia magna* performed with an increased amount of Z-Isomer (isomer ratio (E/Z) = 65/35 demonstrated an at least comparable, potentially lower ecotoxicological profile than the parent E-isomer, demonstrating that there is no further risk for the aquatic compartment (please refer to CA 8.2.4.1 M-030533-01-1). Taking this information into account both isomers can be evaluated as sum of E+Z-isomers, providing a conservative environmental risk assessments.

Any distribution or reproduction of this document for regulatory purposes without the consent of the respective affiliate constitutes a violation of the applicable license agreement.



Table CP 9- 2: Active substance and degradation products addressed in this document

Compound / Codes	Chemical Structure	Explanation for Consideration	Considered for
<b>Fluoxastrobin</b> (HEC 5725)  <b>HEC 5725-Z-Isomer</b>	<p><i>E</i>-isomer</p>  <p><i>Z</i>-isomer</p> 	<p>active substance</p> <p>photolytic metabolite</p>	<p>PEC<sub>soil</sub> PEC<sub>gw</sub> PEC<sub>sw</sub> &amp; PEC<sub>sed</sub> As a worst case approach, the sum of both isomers (Fluoxastrobin E-Z Isomers) is considered for exposure and risk assessment</p>
<b>HEC 5725-carboxylic acid</b> (HEC 7180, M40)			<p>PEC<sub>soil</sub> PEC<sub>gw</sub> PEC<sub>sw</sub> &amp; PEC<sub>sed</sub></p>
<b>HEC 5725-E-des-chlorophenyl</b> (HEC 7155, M48)		<p>occurrence in aerobic soil (&gt;10%) - water/sediment study (&gt;10% in water)</p>	<p>PEC<sub>soil</sub> PEC<sub>gw</sub> PEC<sub>sw</sub> &amp; PEC<sub>sed</sub></p>
<b>2-chlorophenol</b> (M82)		<p>occurrence in aerobic soil (&gt;10%)</p>	<p>PEC<sub>soil</sub> PEC<sub>gw</sub> PEC<sub>sw</sub> &amp; PEC<sub>sed</sub></p>

**Definition of the residue for risk assessment**

For Details please refer to MCA 7.

This document is copyrighted by Bayer AG (or its respective affiliates) and its content for regulatory purposes is prohibited and constitutes a violation of the underlying license agreement.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9- 3: Definition of the residue for risk assessment

Compartment	Residue Definition for Risk Assessment
Soil	fluoxastrobin ( <i>E</i> - isomer), HEC 5725 - <i>Z</i> -isomer, HEC 5725-carboxylic acid ( <i>M40</i> ), HEC 5725- <i>E</i> -des-chlorophenyl ( <i>M48-E</i> ), 2-chlorophenol ( <i>M82</i> )
Groundwater	fluoxastrobin ( <i>E</i> -isomer), HEC 5725- <i>Z</i> -isomer, HEC 5725-carboxylic acid ( <i>M40</i> ), HEC 5725- <i>E</i> -des-chlorophenyl ( <i>M48-E</i> ), 2-chlorophenol ( <i>M82</i> )
Surface water	fluoxastrobin ( <i>E</i> - isomer), HEC 5725- <i>Z</i> -isomer, HEC 5725-carboxylic acid ( <i>M40</i> ), HEC 5725- <i>E</i> -des-chlorophenyl ( <i>M48-E</i> )
Sediment	fluoxastrobin ( <i>E</i> - isomer), HEC 5725- <i>Z</i> -isomer
Air	none


This document is copyright protected.  
Any distribution, reproduction or publication requires  
the consent of Bayer AG (or its respective affiliate).  
Any use of the document or its content for regulatory or  
any other commercial purpose is prohibited and constitutes  
a violation of the underlying license agreement.

**CP 9.1 Fate and behaviour in soil**

For detailed information on the fate and behaviour in soil please refer to MCA Section 7, data point 7.1.

The proposed degradation pathway of fluoxastrobin in soil is shown in Figure CP 9.1- 1.

**Figure CP 9.1- 1: Proposed degradation pathway of fluoxastrobin in soil (major degradation products).**



**CP 9.1.1 Rate of degradation in soil**

No specific studies with the formulation are required. For further information on the fate and behaviour in soil please refer to MCA Section 7, data points 7.1.1 and 7.1.2.

**CP 9.1.1.1 Laboratory studies**

For information on laboratory studies please refer to MCA Section 7, data point 7.1.2.1.

**CP 9.1.1.2 Field studies**

For information on field studies please refer to MCA Section 7, data point 7.1.2.2.





#### CP 9.1.1.2.1 Soil dissipation studies

For information on field dissipation studies please refer to MCA Section 7, data point 7.1.2.2.1.

#### CP 9.1.1.2.2 Soil accumulation studies

For information on field accumulation studies please refer to MCA Section 7, data point 7.1.2.2.2.

#### CP 9.1.2 Mobility in the soil

For information on mobility studies please refer to MCA Section 7, data point 7.1.4.

##### CP 9.1.2.1 Laboratory studies

For information on laboratory studies please refer to MCA Section 7, data point 7.1.4.1.

##### CP 9.1.2.2 Lysimeter studies

For information on lysimeter studies please refer to MCA Section 7, data point 7.1.4.2.

##### CP 9.1.2.3 Field leaching studies

For information on field leaching studies please refer to MCA Section 7, data point 7.1.4.3.

*This document is copyright protected. Any distribution, reproduction or publication requires the consent of Bayer AG (or its respective affiliates) and constitutes any other commercial purpose is prohibited and constitutes a violation of the underlying license agreement.*



**CP 9.1.3 Estimation of concentrations in soil**

New calculations were performed to reflect findings from new studies presented in the active substance dossier, section 7 “Fate and behaviour in the environment”. In addition these calculations considered the most recent guidance documents for exposure calculations. Calculations of predicted environmental concentrations in soil (PEC<sub>soil</sub>) are presented below.

**Predicted environmental concentrations in soil (PECs)**

**Endpoints for PEC<sub>soil</sub>**

For deriving the respective end points please refer to MCA Section 7, data point 7.

**Table CP 9.1.3- 1: Key modelling input parameters for fluoxastrobin and its metabolites**

Compound	Worst case DT <sub>50</sub> non-normalised [days]	Maximum occurrence in soil [%]	Molar mass [g/mol]	Molar mass correction factor	
Fluoxastrobin (E+Z)	DFOP: k <sub>1</sub> fast 0.01741 1/d, k <sub>2</sub> slow 0.002913 1/d, g <sub>fast</sub> 0.4996 (rates equivalent to: DT <sub>50</sub> fast phase 39.81 d, DT <sub>50</sub> slow phase 23 <sup>4</sup> d, g <sub>slow</sub> 0.4996)	DFOP: DT <sub>50</sub> initial 86.41 d <sup>1</sup> , DT <sub>90</sub> initial 552.8 d <sup>2</sup>	100	458.8	1
HEC 5725-E-des-chlorophenyl	9.57 <sup>2,5</sup>	32.2	348.3	0.7592	
HEC 5725-carboxylic acid	28.6 <sup>3,6</sup>	16.9	417.8	0.9106	
2-chlorophenol	23 <sup>4</sup>	49.2 <sup>7</sup>	128.56	0.2802	

1: worst case non-normalized field site (Hurston R812404) with worst-case DFOP DT<sub>90, initial</sub> value

2: worst case non-normalized apparent field decline DT<sub>90</sub> value.

3: worst case non-normalized laboratory DT<sub>50</sub> value.

4: worst case DT<sub>50</sub> value according to the recommendations of EFSA (EFSA, 2007)

5: [REDACTED], 2015; M-534457-01-1 (see MCA 7.1.2.2.1)

6: [REDACTED], 2015; M-534569-01-1 (see MCA 7.1.2.1.2)

7: theoretical estimation by EFSA (EFSA, 2007)

Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

**Report:** KCP 9.1.3/01 [REDACTED]; [REDACTED]; 2015; M-537905-01-1  
**Title:** Fluoxastrobin (FXA) and metabolites: PEC<sub>soil</sub> EUR - Use in cereals and onions in Europe  
**Report No.:** EnSa-15-0541  
**Document No.:** M-537905-01-1  
**Guideline(s):** not applicable  
**Guideline deviation(s):** not applicable  
**GLP/GEP:** no

**Methods and Materials:** The predicted environmental concentrations in soil (PEC<sub>soil</sub>) of fluoxastrobin and its metabolites were estimated based on a first tier approach using a Microsoft® Excel spreadsheet. A bulk density of 1.5 kg/L and a soil mixing depths of 5 cm were used as recommended by FOCUS (1996) and EU Commission (1995, 2000). The accumulation potential of fluoxastrobin after long term use was also assessed, employing the mixing depth of 26 cm for the calculation of the background concentration.

Detailed application data used for simulation of PEC<sub>soil</sub> were compiled in Table CP 9.1.3- 2.

**Table CP 9.1.3- 2: Application pattern used for PEC<sub>soil</sub> calculations of fluoxastrobin**

Individual crop	FOCUS crop used for interception	Application				Amount reaching soil per season application [g a.s./ha]
		Rate per season [g a.s./ha]	Interval [days]	Plant interception [%]	BBCH stage	
Cereals	Cereals	2 × 150	14	2 × 80	2 × 30-69	2 × 30.0
Cereals	Cereals	2 × 125	14	2 × 80	2 × 30-61	2 × 25.0
Onions	Onions	125	10	2 × 10	2 × 15-47	2 × 112.5

**Substance Specific Parameters:** The compound specific input parameters (endpoints for PEC<sub>soil</sub> calculations) are summarized in Table CP 9.1.3- 1.

**Findings:** The maximum PEC<sub>soil</sub> values for fluoxastrobin and its metabolites are summarised in Table CP 9.1.3- 3. The maximum, short-term and long-term PEC<sub>soil</sub> values and the time weighted average values (TWA<sub>soil</sub>) are provided in tables 9.1.3-4 and 9.1.3-5.

**Table CP 9.1.3- 3: Maximum PEC<sub>soil</sub> of fluoxastrobin and its metabolites for the uses assessed**

Use Pattern	Fluoxastrobin (E+Z) PEC <sub>soil</sub> [mg/kg]	HEC 5725-E-des-chlorophenyl PEC <sub>soil</sub> [mg/kg]	HEC 5725-carboxylic acid PEC <sub>soil</sub> [mg/kg]	2-chlorophenol PEC <sub>soil</sub> [mg/kg]
Cereals 2×150 g a.s./ha, 14 days, 2×80%	0.075	0.019	0.011	0.009
Cereals 2×125 g a.s./ha, 14 days, 2×80%	0.062	0.016	0.009	0.008
Onions 2×125 g a.s./ha, 10 days, 2×10%	0.286	0.071	0.041	0.036



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.1.3- 4: Cereals, 2 × 150 g a.s./ha: PEC<sub>soil</sub> (actual) of fluoxastrobin and its metabolites

	Time [days]	Cereals			
		2 × 150 g a.s./ha, 14 days app. interval, 2 × 80% interception			
		Fluoxastrobin (E+Z)	HEC 5725-E-des-chlorophenyl	HEC 5725-carboxylic acid	2-chlorophenol
		PEC <sub>soil</sub> [mg/kg]	PEC <sub>soil</sub> [mg/kg]	PEC <sub>soil</sub> [mg/kg]	PEC <sub>soil</sub> [mg/kg]
Initial	0	0.075	0.019	0.010	0.009
Short term	1	0.074	0.018	0.010	0.009
	2	0.073	0.018	0.010	0.009
	4	0.072	0.018	0.010	0.008
Long term	7	0.070	0.018	0.009	0.007
	14	0.066	0.017	0.008	0.006
	21	0.062	0.016	0.006	0.005
	28	0.058	0.015	0.005	0.004
	42	0.052	0.014	0.004	0.003
	50	0.049	0.013	0.003	0.002
	100	0.036	0.009	0.001	<0.001

Table CP 9.1.3- 5: Cereals, 2 × 150 g a.s./ha: TWAC<sub>soil</sub> of fluoxastrobin and its metabolites

	Time [days]	Cereals			
		2 × 150 g a.s./ha, 14 days app. interval, 2 × 80% interception			
		Fluoxastrobin (E+Z)	HEC 5725-E-des-chlorophenyl	HEC 5725-carboxylic acid	2-chlorophenol
		TWAC <sub>soil</sub> [mg/kg]	TWAC <sub>soil</sub> [mg/kg]	TWAC <sub>soil</sub> [mg/kg]	TWAC <sub>soil</sub> [mg/kg]
Initial	0	---	---	---	---
Short term	1	0.073	0.019	0.010	0.009
	2	0.074	0.018	0.010	0.009
	4	0.073	0.018	0.010	0.009
Long term	7	0.072	0.018	0.010	0.008
	14	0.070	0.018	0.009	0.007
	21	0.068	0.017	0.008	0.007
	28	0.066	0.017	0.008	0.006
	42	0.062	0.016	0.007	0.005
	50	0.060	0.016	0.006	0.005
	100	0.051	0.013	0.004	0.003

This document is copyright protected. Any distribution or reproduction of its content for regulatory and commercial purposes without the consent of Bayer AG or its respective affiliates is prohibited and constitutes a violation of the underlying license agreement.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.1.3- 6: Cereals, 2 × 125 g a.s./ha: PEC<sub>soil</sub> (actual) of fluoxastrobin and its metabolites

	Time [days]	Cereals			
		2 × 125 g a.s./ha, 14 days app. interval, 2 × 80% interception			
		Fluoxastrobin (E+Z)	HEC 5725-E-des-chlorophenyl	HEC 5725-carboxylic acid	2-chlorophenol
		PEC <sub>soil</sub> [mg/kg]	PEC <sub>soil</sub> [mg/kg]	PEC <sub>soil</sub> [mg/kg]	PEC <sub>soil</sub> [mg/kg]
Initial	0	0.062	0.016	0.009	0.008
Short term	1	0.062	0.015	0.009	0.007
	2	0.061	0.015	0.008	0.007
	4	0.060	0.015	0.008	0.007
Long term	7	0.058	0.015	0.007	0.006
	14	0.055	0.014	0.006	0.005
	21	0.051	0.013	0.005	0.004
	28	0.048	0.013	0.004	0.003
	42	0.043	0.011	0.003	0.002
	50	0.041	0.011	0.003	0.002
	100	0.030	0.008	0.001	<0.001

Table CP 9.1.3- 7: Cereals, 2 × 125 g a.s./ha: TWAC<sub>soil</sub> of fluoxastrobin and its metabolites

	Time [days]	Cereals			
		2 × 125 g a.s./ha, 14 days app. interval, 2 × 80% interception			
		Fluoxastrobin (E+Z)	HEC 5725-E-des-chlorophenyl	HEC 5725-carboxylic acid	2-chlorophenol
		TWAC <sub>soil</sub> [mg/kg]	TWAC <sub>soil</sub> [mg/kg]	TWAC <sub>soil</sub> [mg/kg]	TWAC <sub>soil</sub> [mg/kg]
Initial	0	---	---	---	---
Short term	1	0.062	0.015	0.009	0.007
	2	0.062	0.015	0.009	0.007
	4	0.061	0.015	0.008	0.007
Long term	7	0.060	0.015	0.008	0.007
	14	0.058	0.015	0.007	0.006
	21	0.057	0.014	0.007	0.006
	28	0.055	0.014	0.006	0.005
	42	0.052	0.013	0.006	0.004
	50	0.050	0.013	0.005	0.004
	100	0.042	0.011	0.003	0.002

This document is copyright protected. Any distribution or reproduction of its content for regulatory and commercial purposes without the consent of Bayer AG or its respective affiliates is prohibited and constitutes a violation of the underlying license agreement.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.1.3- 8: Onions, 2 × 125 g a.s./ha: PEC<sub>soil</sub> (actual) of fluoxastrobin and its metabolites

	Time [days]	Onions 2 × 125 g a.s./ha, 10 days app. interval, 2 × 10% interception			
		Fluoxastrobin (E+Z)	HEC 5725-E-des-chlorophenyl	HEC 5725-carboxylic acid	2-chlorophenol
		PEC <sub>soil</sub> [mg/kg]	PEC <sub>soil</sub> [mg/kg]	PEC <sub>soil</sub> [mg/kg]	PEC <sub>soil</sub> [mg/kg]
Initial	0	0.286	0.071	0.041	0.036
Short term	1	0.283	0.070	0.040	0.035
	2	0.280	0.070	0.039	0.034
	4	0.275	0.069	0.037	0.032
Long term	7	0.267	0.067	0.035	0.029
	14	0.250	0.064	0.029	0.024
	21	0.235	0.061	0.025	0.019
	28	0.221	0.058	0.021	0.015
	42	0.197	0.052	0.015	0.010
	50	0.186	0.049	0.014	0.008
	100	0.135	0.034	0.004	0.002

Table CP 9.1.3- 9: Onions, 2 × 125 g a.s./ha: TWAC<sub>soil</sub> of fluoxastrobin and its metabolites

	Time [days]	Onions 2 × 125 g a.s./ha, 10 days app. interval, 2 × 10% interception			
		Fluoxastrobin (E+Z)	HEC 5725-E-des-chlorophenyl	HEC 5725-carboxylic acid	2-chlorophenol
		TWAC <sub>soil</sub> [mg/kg]	TWAC <sub>soil</sub> [mg/kg]	TWAC <sub>soil</sub> [mg/kg]	TWAC <sub>soil</sub> [mg/kg]
Initial	0	---	---	---	---
Short term	1	0.284	0.071	0.041	0.035
	2	0.283	0.070	0.040	0.035
	4	0.280	0.070	0.039	0.034
Long term	7	0.276	0.069	0.038	0.032
	14	0.267	0.067	0.035	0.029
	21	0.259	0.066	0.032	0.027
	28	0.251	0.064	0.030	0.024
	42	0.237	0.061	0.026	0.020
	50	0.230	0.059	0.024	0.019
	100	0.194	0.050	0.016	0.011

This document is copyright protected. Any distribution or reproduction of its content for regulatory and commercial purposes without the consent of Bayer AG or its respective affiliates is prohibited and constitutes a violation of the underlying license agreement.



**Potential accumulation in soil:**

The accumulation potential after long term use was also assessed. The results for a standard-mixing depth of 20 cm for an arable crop with tillage are presented in Table CP 9.1.3- 10.

**Table CP 9.1.3- 10: PEC<sub>soil</sub> of fluoxastrobin taking the effect of accumulation into account (mixing depth of 20 cm)**

Use Pattern	PEC <sub>soil</sub>	Fluoxastrobin (E+Z)
		[mg/kg]
Cereals 2×150 g a.s./ha, 14 days, 2×80%	plateau	0.005
	total*	0.080
Cereals 2×125 g a.s./ha, 14 days, 2×80%	plateau	0.005
	total*	0.067
Onions 2×125 g a.s./ha, 10 days, 2×10%	plateau	0.020
	total*	0.306

\* total = plateau (background concentration after multi-year use) + max. PEC<sub>soil</sub>


This document is copyright protected. Any distribution, reproduction or publication requires the consent of Bayer AG (or its respective regulatory or constituting entities). Any use of the document or its content for regulatory or constituting purposes without the prior written consent of Bayer AG constitutes a violation of the underlying license agreement.

**CP 9.2 Fate and behaviour in water and sediment**

The proposed degradation pathway of fluoxastrobin in water and sediment is shown in Figure CP 9.2-1.

For information on the fate and behaviour in water and sediment please refer to MCA Section 7, data point 7.2.

**Figure CP 9.2- 1: Proposed bio-degradation pathway of fluoxastrobin in water and sediment (major degradation products)**



Bound residues  
(max. 12.7% aerobic; 36.2% anaerobic)

14C-CO2  
(max. 2.9% aerobic)





**CP 9.2.1 Aerobic mineralisation in surface water**

For information on aerobic mineralisation in surface water studies please refer to MCA Section 7, data point 7.2.2.2.

**CP 9.2.2 Water/sediment study**

For information on water/sediment studies please refer to MCA Section 7, data point 7.2.2.3.

**CP 9.2.3 Irradiated water/sediment study**

For information on irradiated water/sediment studies please refer to MCA Section 7, data point 7.2.2.4.

*This document is copyright protected.  
Any distribution, reproduction or publication requires  
the consent of Bayer AG (or its respective affiliate).  
Any use of the document or its content for regulatory or  
any other commercial purpose is prohibited and constitutes  
a violation of the underlying license agreement.*



**CP 9.2.4 Estimation of concentrations in groundwater**

Calculations were performed, to reflect findings from new studies presented in the active substance dossier, section 7 “Fate and behaviour in the environment”. In addition these calculations consider the most recent guidance documents for exposure calculations.

Calculations of predicted environmental concentrations in groundwater (PEC<sub>gw</sub>) are presented below.

**PEC<sub>gw</sub> modelling approach**

The predicted environmental concentrations in groundwater (PEC<sub>gw</sub>) for the active substance were calculated using the simulation models FOCUS PEARL and FOCUS PELMO following the recommendations of the FOCUS working group on groundwater scenarios. Further, where a crop of interest is defined for Châteaudun scenario, FOCUS MACRO simulations were performed (EFSA Guidance Document, 2014<sup>1</sup>).

The leaching calculations were run over 26 years, as proposed for pesticides which may be applied every year. The first six years are a ‘warm up’ period; only the last 20 years were considered for the assessment of the leaching potential. The 80<sup>th</sup> percentile of the mean annual groundwater concentrations in the percolate at 1 m depth under a treated plantation were evaluated and were taken as the relevant PEC<sub>GW</sub> values. In respect to the assessment of a potential groundwater contamination this shallow depth reflects a worst case. The effective long-term groundwater concentrations will be even lower due to dilution in the upper groundwater layer.

Crop interception will reduce the amount of a compound reaching the soil and therefore this was taken into account depending on the growth stage at application. The interception rates follow the EFSA Guidance Document (2014)<sup>1</sup> recommendations (Table CP 9.2.4- 1).

**Table CP 9.2.4- 1: EFSA (2014) groundwater crop interception values**

Crop	Crop stage Interception [%]						
	Bare emergence	Leaf development	Stem elongation		Flowering		Senescence Ripening
	00 - 09	10 - 19	20 - 29	30 - 39	40 - 69	70 - 89	90 - 99
Winter cereals	0	0	20 (tillering)	80 (elong.)	90	80	80
Spring cereals	0	0	20 (tillering)	80 (elong.)	90	80	80
Onions		10	20 - 39		40 - 89		60
			25		40		

**Endpoints for PEC<sub>gw</sub>**

For deriving the respective endpoints please refer to MCA7, data point 7.1.

<sup>1</sup> EFSA (2014): EFSA Guidance Document for evaluating laboratory and field dissipation studies to obtain DegT50 values of active substances of plant protection products and transformation products of these active substances in soil. EFSA Journal 2014;12(5):3662.



Table CP 9.2.4- 2: Key modelling input parameters for fluoxastrobin and its metabolites

Compound	DT <sub>50</sub> soil [days]	Koc [mL/g]	Kom [mL/g]	FREUNDLICH exponent 1/n
Fluoxastrobin (E+Z)	38.89	752.0	436.2	0.8584
HEC 5725-E-des-chlorophenyl	56.7	19.3 <sup>1)</sup>	11.2 <sup>1)</sup>	0.9367 <sup>2)</sup>
HEC 5725-carboxylic acid	17.01	56.4	32.8	0.9043
2-chlorophenol	23.0	104.7	60	0.8520

- 1) geomean of neutral pH cluster
- 2) Arithm. mean of neutral pH cluster

CP 9.2.4.1 Calculation of concentrations in groundwater

Predicted environmental concentrations in groundwater (PEC<sub>gw</sub>)

*PEC<sub>gw</sub> values for the use in cereals and onions - FOCUS PEARL and PELMO*

**Report:** KCP 9.2.4.1/01 [redacted]; 2015; M-537900-01-1  
**Title:** Fluoxastrobin (FXA) and metabolites: PEC<sub>gw</sub> FOCUS PEARL, PELMO EUR - Use in cereals in Europe  
**Report No.:** Ensa-15-0545  
**Document No.:** M-537900-01-1  
**Guideline(s):** not applicable  
**Guideline deviation(s):** not applicable  
**GLP/GEP:** no

**Report:** KCP 9.2.4.1/02 [redacted]; 2015; M-537902-01-1  
**Title:** Fluoxastrobin (FXA) and metabolites: PEC<sub>gw</sub> FOCUS PEARL, PELMO EUR - Use in Onions in Europe  
**Report No.:** Ensa-15-0551  
**Document No.:** M-537902-01-1  
**Guideline(s):** not applicable  
**Guideline deviation(s):** not applicable  
**GLP/GEP:** no

The predicted environmental concentrations in groundwater (PEC<sub>gw</sub>) for fluoxastrobin and its metabolites were calculated using the simulation model FOCUS PEARL (version 4.4.4) and FOCUS PELMO (version 5.53). Crop interception was taken into account according to the BBCH growth stage, as recommended by EFSA (EFSA (2014), FOCUS (2014)). The absolute dates for applications based on BBCH codes given in the GAP were determined using AppDate2 (Klein (2010)), a German regulatory tool for estimating application dates and crop interception.

Typically, a leaching assessment is carried out considering aerobic conditions as a common agricultural situation. Therefore, observed major aerobic metabolites were taken into account, implementing their amounts and behaviour as observed under aerobic conditions.

However, in anaerobic soil, a further fast degrading major metabolite, HEC5725-carboxylic acid (HEC7180, M40), was identified (16.9 % at day 120), which did not occur under aerobic conditions. Based on these observations, a conservative anaerobic leaching assessment was carried out for this metabolite, respectively.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Anaerobic leaching scenario:

Under common agricultural situations in Europe, considering e.g. climatic conditions or slope of fields, it is obviously unrealistic, that a total treated agricultural field or area turns anaerobic, each year after application and lasting for a long time period, as typically considered for aerobic leaching assessments. Such conditions would make farming effectively impossible.

Therefore, two more realistic, but still very conservative scenarios were considered here:

**Scenario 1:** Anaerobic conditions may occur regularly in plane fields or cropping areas, when rain water remains in small sinks and furrows with low permeability. In this case, only a relatively small percentage of the total cropped area or field would be affected.

**Scenario 2:** Anaerobic conditions on larger scale may occur due to flooding along rivers. Typically, this flooding will not occur regularly or each year, only with large time intervals in between.

The following assumptions were made to address these two scenarios. Partly, additional safety factors were applied to address uncertainties in the estimation.

Here, it is implicitly included that anaerobic conditions occur more or less immediately after application (1 day later) and that anaerobic conditions are as strict as simulated in the lab. In reality, it may take considerable time after ponding until anaerobic conditions occur, because the remaining oxygen in soil and water has to be consumed by microbes, first. Furthermore, in the lab studies anaerobic conditions are ensured by ventilating the samples with nitrogen. Such conditions will not appear in reality.

Therefore, it has to be noted, that the described assumptions and scenarios are highly conservative.

**Table CP 9.2.4.1- 1: Assumptions used for anaerobic leaching scenarios**

Scenario	Assumption	Safety factor	actually used
1	not more than 10 % area of an agricultural field becomes anaerobic, every year shortly after application	1	<b>application rate</b> reduced to <b>10 %</b> , applied <b>every year</b> (application rate 100 %, applied every year, PEC <sub>gw</sub> divided by 10)
2	Calculation base for dimension of levees, dykes and flood plains along rivers are 100-year-floodings. Hence, ponding on larger areas can be assumed to occur in average every 100 years	10	<b>application rate 100 %</b> , applied <b>every 10 years</b>
both	Farmer will not apply on saturated and ponded fields. Therefore, it is assumed, that parent compound degrades 1 day under aerobic conditions before anaerobic conditions occur.		<u>degradation</u> time for <u>parent</u> before anaerobic = <u>1 day</u>
both	Anaerobic conditions usually will not last for longer than 1 week. Maximum occurrence of metabolite might not yet be reached at this time.		<u>maximum occurrence</u> in anaerobic soil of M40 = <u>16.9%</u> (found after 120 d)
both	After an anaerobic period, normal aerobic agricultural conditions may dominate in soil again. Thus, aerobic degradation of the anaerobic metabolite is assessed.		<u>Aerobic lab DT<sub>50</sub></u> of <u>17.01 d</u> (M40)

Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) GPseudo application of anaerobic metabolite:

The anaerobic metabolite is assumed to be applied directly to the soil by pseudo application. Hence, no “pathway”-calculation was done in which the parent is applied. This is considered the only plausible but conservative way to account for the anaerobic formation (expressed by the maximum occurrence) and the aerobic degradation of the anaerobic metabolite. Applying the aerobic pathway for groundwater calculations may disregard the formation under anaerobic conditions.

Detailed application data used for simulation of  $PEC_{gw}$  for all compounds were compiled in Table CP 9.2.4.1- 2.

Table CP 9.2.4.1- 2: Application pattern used for  $PEC_{gw}$  calculations

Individual crop	FOCUS crop used for interception	Application			Amount reaching soil per season application [g a.s./ha]
		Rate per season [g a.s./ha]	Interval [days]	Plant interception [%]	
Winter & spring cereals, GAP	-	2 × 150	14	-	-
Spring cereals 1, simulation	Spring cereals	2 × 150	14	2 × 80	2 × 30.0
Spring cereals 2, simulation <sup>2)</sup>	Spring cereals	2 × 22.68 <sup>1)</sup>	14	2 × 80	2 × 4.54 <sup>1)</sup>
Winter cereals 1, simulation	Winter cereals	2 × 150	14	2 × 80	2 × 30.0
Winter cereals 2, simulation <sup>2)</sup>	Winter cereals	2 × 22.68 <sup>1)</sup>	14	2 × 80	2 × 4.54 <sup>1)</sup>
Winter & spring cereals, GAP	-	2 × 125	14	-	-
Spring cereals 3, simulation	Spring cereals	2 × 125	14	2 × 80	2 × 25.0
Spring cereals 4, simulation <sup>2)</sup>	Spring cereals	2 × 18.90 <sup>1)</sup>	14	2 × 80	2 × 3.78 <sup>1)</sup>
Winter cereals 3, simulation	Winter cereals	2 × 125	14	2 × 80	2 × 25.0
Winter cereals 4, simulation <sup>2)</sup>	Winter cereals	2 × 18.90 <sup>1)</sup>	14	2 × 80	2 × 3.78 <sup>1)</sup>
Onions, GAP	-	2 × 125	10	-	-
Onions 1, simulation	Onions	2 × 125	10	2 × 10	2 × 112.5
Onions 2, simulation <sup>2)</sup>	Onions	2 × 18.90 <sup>1)</sup>	10	2 × 10	2 × 17.0 <sup>1)</sup>

<sup>1)</sup> Pseudo application [g metabolite /ha]

<sup>2)</sup> Pseudo application pattern for anaerobic metabolite HEC 5725-carboxylic acid: parent rate – 1 d degradation, corrected for molar masses and maximum occurrence in anaerobic soil (= 100% metabolite rate)

For cereal and onion applications, absolute dates were derived for the simulation runs. All application dates are summarised in the table below.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.4.1- 3: Application dates and related information for fluoxastrobin as used for the simulation runs

Individual crop	Spring cereals 1 – 4	Winter cereals 1 – 4	Onions 1 – 2
Repeat Interval for App. Events	Every Year	Every Year	Every Year
Application Technique	Spray	Spray	Spray
Absolute / Relative to	Absolute	Absolute	Absolute
Scenario	1 <sup>st</sup> App. Date (Julian day) Offset	1 <sup>st</sup> App. Date (Julian day) Offset	1 <sup>st</sup> App. Date (Julian day) Offset
Chateaudun	10 Apr (100)	21 Apr (119)	29 May (149)
Hamburg	28 Apr (118)	09 Apr (109)	29 May (149)
Jokioinen	05 Jun (156)	25 May (145)	06 Jun (159)
Kremsmuenster	28 Apr (118)	19 Apr (109)	29 May (149)
Okehampton	2 Apr (112)	15 Apr (105)	-
Piacenza	-	10 Apr (100)	-
Porto	16 Apr (106)	30 Mar (89)	08 Apr (98)
Sevilla	-	06 Jan (6)	-
Thjva	-	02 Mar (61)	14 May (134)

Substance specific and model related input parameters for FOCUS PEARL & PELMO PEC<sub>gw</sub> calculations are summarised in Table CP 9.2.4.1- 4. Degradation pathway related parameters are given in Table CP 9.2.4.1- 5.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.4.1- 4: Compound input parameters for fluoxastrobin and its metabolites

Parameter	Unit	Fluoxastrobin (E+Z)	HEC 5725-E-des-chlorophenyl	HEC 5725-carboxylic acid	2-chlorophenol
<b>Common</b>					
Molar Mass	[g/mol]	458.8	348.3	417.8	128.56 (264.8 <sup>1)</sup> )
Solubility	[mg/L]	2.292	9600	244 000	23 000
Vapour Pressure	[Pa]	5.63E-10	6.00E-05	7.00E-04	1.44E+02
Freundlich Exponent		0.8584	0.9367	0.9043	0.8520
Plant Uptake Factor		0.0	0.0	0.0	0.0
Walker Exponent		0.7	0.7	0.7	0.7
<b>PEARL Parameters</b>					
Substance Code		FXA	E-des	Carb	Chlph
DT <sub>50</sub>	[days]	38.89	56.7	7.01	7.0
Molar Activ. Energy	[kJ/mol]	65.4	65.4	65.4	65.4
K <sub>om</sub>	[mL/g]	436.2	11.2	32.8	60.7
K <sub>f</sub>	[mL/g]	-	-	-	-
<b>PELMO Parameters</b>					
Substance Code		AS	A1	AS	B1
Rate Constant	[1/day]	0.0179	0.0122	0.0407	0.03014
Q <sub>10</sub>		2.58	2.58	2.58	2.58
K <sub>oc</sub>	[mL/g]	452.0	29.3	56.4	104.7

1) PELMO parameters: An auxiliary molar mass of 2-chlorophenol is introduced, to compensate for the low split degradation rate and to ensure the correct mass flux.

Table CP 9.2.4.1- 5: Degradation pathway related parameters for fluoxastrobin and its metabolites

Degradation fraction from → to (FOCUS PEARL)	1 FXA → Chlph 0.5145 FXA → E-des
Degradation rate from → to (FOCUS PELMO)	0.00917 Active Substance → A1 0.0086 Active Substance → B1 0.0122 A1 →  O2 0.03014 B1 →  CO2

**Findings:** PEC<sub>gw</sub> were evaluated as the 80<sup>th</sup> percentile of the mean annual leachate concentration at 1 m soil depth. FOCUS PEARL and PELMO PEC<sub>gw</sub> results for fluoxastrobin and its metabolites after application to winter and spring cereals and onions are given in Table CP 9.2.4.1- 6.



Table CP 9.2.4.1- 6: Spring cereals: FOCUS PEARL & PELMO PEC<sub>gw</sub> results of fluoxastrobin and its metabolites

Use Pattern	Spring cereals 1 - 3, 2 × 150 g a.s./ha, 2 × 80% interception, 14 d interval			
	Fluoxastrobin (E+Z)	HEC 5725-E- des- chlorophenyl	2-chlorophenol	HEC 5725- carboxylic acid <sup>1)</sup>
FOCUS PEARL	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Chateaudun	<0.001	1.093	<0.001	<0.001
Hamburg	<0.001	3.086	<0.001	<0.001
Jokioinen	<0.001	2.272	<0.001	<0.001
Kremsmuenster	<0.001	1.678	<0.001	<0.001
Okehampton	<0.001	1.678	<0.001	<0.001
Porto	<0.001	1.140	<0.001	<0.001
FOCUS PELMO	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Chateaudun	<0.001	0.903	<0.001	<0.001
Hamburg	<0.001	2.438	<0.001	<0.001
Jokioinen	<0.001	2.213	<0.001	<0.001
Kremsmuenster	<0.001	0.673	<0.001	<0.001
Okehampton	<0.001	1.602	<0.001	<0.001
Porto	<0.001	1.094	<0.001	<0.001

<sup>1)</sup> Pseudo application pattern for the anaerobic metabolite HEC 5725-carboxylic acid (Scenario 1).

This document is copyright protected. Any distribution, reproduction or its use without the consent of Bayer AG (or its respective affiliates) for regulatory or commercial purposes is prohibited and constitutes a violation of the underlying license agreement.





Table CP 9.2.4.1- 7: Winter cereals: FOCUS PEARL & PELMO PEC<sub>gw</sub> results of fluoxastrobin and its metabolites

Use Pattern	Winter cereals 1 - 3, 2 × 150 g a.s./ha, 2 × 80% interception, 14 d interval			
	Fluoxastrobin (E+Z)	HEC 5725-E- des- chlorophenyl	2-chlorophenol	HEC 5725- carboxylic acid <sup>1)</sup>
FOCUS PEARL	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Chateaudun	<0.001	1.209	<0.001	<0.001
Hamburg	<0.001	2.478	<0.001	<0.001
Jokioinen	<0.001	2.688	<0.001	<0.001
Kremsmuenster	<0.001	1.561	<0.001	<0.001
Okehampton	<0.001	1.571	<0.001	<0.001
Piacenza	<0.001	0.996	<0.001	<0.001
Porto	<0.001	0.939	<0.001	<0.001
Sevilla	<0.001	0.276	<0.001	<0.001
Thiva	<0.001	0.831	<0.001	<0.001
FOCUS PELMO	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Chateaudun	<0.001	0.075	<0.001	<0.001
Hamburg	<0.001	2.566	<0.001	<0.001
Jokioinen	<0.001	2.767	<0.001	<0.001
Kremsmuenster	<0.001	1.720	<0.001	<0.001
Okehampton	<0.001	1.636	<0.001	<0.001
Piacenza	<0.001	1.286	<0.001	<0.001
Porto	<0.001	1.118	<0.001	<0.001
Sevilla	<0.001	0.332	<0.001	<0.001
Thiva	<0.001	0.507	<0.001	<0.001

<sup>1)</sup> Pseudo application pattern for the anaerobic metabolite HEC 5725-carboxylic acid (Scenario 1).

Table CP 9.2.4.1- 8: Spring cereals: FOCUS PEARL & PELMO PEC<sub>gw</sub> results of fluoxastrobin and its metabolites

Use Pattern	Spring cereals 4 & 5, 2 × 125 g a.s./ha, 2 × 80% interception, 14 d interval			
	Fluoxastrobin (E+Z)	HEC 5725-E- des- chlorophenyl	2-chlorophenol	HEC 5725- carboxylic acid <sup>1)</sup>
FOCUS PEARL	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Chateaudun	<0.001	0.901	<0.001	<0.001
Hamburg	<0.001	2.538	<0.001	<0.001
Jokioinen	<0.001	1.875	<0.001	<0.001
Kremsmuenster	<0.001	1.385	<0.001	<0.001
Okehampton	<0.001	1.386	<0.001	<0.001
Porto	<0.001	0.945	<0.001	<0.001
FOCUS PELMO	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Chateaudun	<0.001	0.744	<0.001	<0.001
Hamburg	<0.001	2.015	<0.001	<0.001
Jokioinen	<0.001	1.829	<0.001	<0.001
Kremsmuenster	<0.001	1.385	<0.001	<0.001
Okehampton	<0.001	1.326	<0.001	<0.001
Porto	<0.001	0.904	<0.001	<0.001

<sup>1)</sup> Pseudo application pattern for the anaerobic metabolite HEC 5725-carboxylic acid (Scenario 1)



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.4.1- 9: Winter cereals: FOCUS PEARL & PELMO PEC<sub>gw</sub> results of fluoxastrobin and its metabolites

Use Pattern	Winter cereals 4 & 5, 2 × 125 g a.s./ha, 2 × 80% interception, 14 d interval			
	Fluoxastrobin (E+Z)	HEC 5725-E-des- chlorophenyl	2-chlorophenol	HEC 5725- carboxylic acid <sup>1)</sup>
FOCUS PEARL	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Chateaudun	<0.001	0.996	<0.001	<0.001
Hamburg	<0.001	2.043	<0.001	<0.001
Jokioinen	<0.001	2.207	<0.001	<0.001
Kremsmuenster	<0.001	1.288	<0.001	<0.001
Okehampton	<0.001	1.300	<0.001	<0.001
Piacenza	<0.001	0.823	<0.001	<0.001
Porto	<0.001	0.793	<0.001	<0.001
Sevilla	<0.001	0.229	<0.001	<0.001
Thiva	<0.001	0.685	<0.001	<0.001
FOCUS PELMO	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Chateaudun	<0.001	0.886	<0.001	<0.001
Hamburg	<0.001	2.106	<0.001	<0.001
Jokioinen	<0.001	2.273	<0.001	<0.001
Kremsmuenster	<0.001	1.421	<0.001	<0.001
Okehampton	<0.001	1.354	<0.001	<0.001
Piacenza	<0.001	1.064	<0.001	<0.001
Porto	<0.001	0.924	<0.001	<0.001
Sevilla	<0.001	0.271	<0.001	<0.001
Thiva	<0.001	0.416	<0.001	<0.001

<sup>1)</sup> Pseudo application pattern for the anaerobic metabolite HEC 5725-carboxylic acid (Scenario 1)

Table CP 9.2.4.1- 10: Onions: FOCUS PEARL & PELMO PEC<sub>gw</sub> results of fluoxastrobin and its metabolites

Use Pattern	Onions 1 - 3, 2 × 125 g a.s./ha, 2 × 10% interception, 10 d interval			
	Fluoxastrobin (E+Z)	HEC 5725-E- des- chlorophenyl	2-chlorophenol	HEC 5725- carboxylic acid <sup>1)</sup>
FOCUS PEARL	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Chateaudun	<0.001	5.101	<0.001	<0.001
Hamburg	<0.001	8.759	<0.001	0.0002
Jokioinen	<0.001	7.786	<0.001	<0.001
Kremsmuenster	<0.001	5.502	<0.001	0.0001
Okehampton	<0.001	2.590	<0.001	<0.001
Porto	<0.001	2.907	<0.001	<0.001
FOCUS PELMO	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Chateaudun	<0.001	4.704	<0.001	<0.001
Hamburg	<0.001	8.457	<0.001	0.0003
Jokioinen	<0.001	7.248	<0.001	<0.001
Kremsmuenster	<0.001	5.783	<0.001	0.0002
Okehampton	<0.001	3.891	<0.001	0.0001
Porto	<0.001	2.174	<0.001	<0.001

<sup>1)</sup> Pseudo application pattern for the anaerobic metabolite HEC 5725-carboxylic acid (Scenario 1).



**Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G**

As described for scenario 1, 100 % of the potential pseudo application rate of anaerobic HEC 5725-carboxylic acid was applied, each year. All  $PEC_{gw}$  values for all groundwater scenarios and application periods resulted already in concentrations  $\leq 0.003 \mu\text{g/L}$ , also without a division by 10. Therefore, a further simulation according Scenario 2, every 10 years, was not carried out anymore, as it is already covered with the first simulation.

**Conclusion:** There are no concerns for groundwater from the use of fluoxastrobin in accordance with the use pattern for the representative formulation.

The concentration of the metabolite HEC 5725-E-des-chlorophenyl (M48) was predicted to reach groundwater at concentrations exceeding  $0.1 \mu\text{g/L}$ . However, the relevance of this metabolite was assessed and the metabolite is non-relevant in groundwater (see Document N4).

*PEC<sub>gw</sub> values for the use in cereals and onions – FOCUS MACRO*

As recommended by FOCUS (2014),  $PEC_{gw}$  were calculated in addition with MACRO 5.5.3, as the Châteaudun scenario has been defined for cereals and onions.

**Report:** KCP 9.2.4.1/03 [redacted]; 2013; M-537903-01-1  
**Title:** Fluoxastrobin (FXA) and metabolites:  $PEC_{gw}$  FOCUS MACRO 5.5.3 EUR - Use in cereals and onions in Europe  
**Report No.:** Ensa-15-0846  
**Document No.:** M-537903-01-1  
**Guideline(s):** not applicable  
**Guideline deviation(s):** not applicable  
**GLP/GEP:** no

The predicted environmental concentrations in groundwater ( $PEC_{gw}$ ) for fluoxastrobin and its metabolites were calculated using the simulation model FOCUS MACRO (version 5.5.3) to simulate macro pore flow for drained soils for Châteaudun scenario. Crop interception was taken into account according to the BBCH growth stage as recommended by EFSA (EFSA (2014), FOCUS (2014)). The absolute dates for applications based on BBCH codes given in the GAP were determined using AppDate2 ([redacted] (2015)), a German regulatory tool for estimating application dates and crop interception.

Typically, a leaching assessment is carried out considering aerobic conditions as a common agricultural situation. Therefore, observed major aerobic metabolites were taken into account, implementing their amounts and behaviour as observed under aerobic conditions. However, in anaerobic soil a further fast degrading major metabolite, HEC5725-carboxylic acid (HEC7180, M40) was identified (6.9 % at day 120), which did not occur under aerobic conditions. Based on these observations, a conservative anaerobic leaching assessment was carried out for this metabolite, respectively.

Anaerobic leaching scenario:

Under common agricultural situations in Europe, considering e.g. climatic conditions or slope of fields, it is obviously unrealistic, that a total treated agricultural field or area turns anaerobic, each year after application and lasting for a long time period, as typically considered for aerobic leaching assessments. Such conditions would make farming effectively impossible.

Therefore, two more realistic, but still very conservative scenarios have been considered here:



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

**Scenario 1:** Anaerobic conditions may occur regularly in plane fields or cropping areas, when rain water remains in small sinks and furrows with low permeability. In this case, only a relatively small percentage of the total cropped area or field would be affected.

**Scenario 2:** Anaerobic conditions on larger scale may occur due to flooding along rivers. Typically, this flooding will not occur regularly or each year, only with large time intervals in between.

The following assumptions have been made to address these two scenarios. Partly, additional safety factors are applied to address uncertainties in the estimation.

Here, it is implicitly included that anaerobic conditions occur more or less immediately after application (1 day later) and that anaerobic conditions are as strict as simulated in the lab. In reality, it may take considerable time after ponding until anaerobic conditions occur, because the remaining oxygen in soil and water has to be consumed by microbes first. Further on, in the lab studies anaerobic conditions are ensured by ventilating the samples with nitrogen. Such conditions will not appear in reality.

Therefore, it has to be noted, that the described assumptions and scenarios are highly conservative.

**Table CP 9.2.4.1- 11: Assumptions used for anaerobic leaching scenarios**

Scenario	Assumption	Safety factor	actually used
1	not more than 10 % area of an agricultural field becomes anaerobic, every year shortly after application		<b>application rate</b> reduced to <b>10 %</b> , applied <b>every year</b> (application rate 100 %, applied every year, $PEC_{gw}$ divided by 10)
2	Calculation base for dimension of levees, dykes and flood plains along rivers are 100-year-floodings. Hence, ponding on larger areas can be assumed to occur in average every 100 years	10	<b>application rate 100 %</b> , applied <b>every 10 years</b>
both	Farmer will not apply on saturated and ponded fields. Therefore it is assumed, that parent compound degrades 1 day under aerobic conditions before anaerobic conditions occur		<u>degradation</u> time for <u>parent</u> before anaerobic = <u>1 day</u>
both	Anaerobic conditions usually will not last for longer than 1 week. Maximum occurrence of metabolite might not yet be reached at this time		<u>maximum occurrence</u> in anaerobic soil of <u>M40</u> = <u>16.9%</u> (found after 120 d)
both	After an anaerobic period, normal aerobic agricultural conditions may dominate in soil again. Thus, aerobic degradation of the anaerobic metabolite is assessed.		<u>Aerobic lab</u> <u>DT<sub>50</sub></u> of <u>17.01 d</u> (M40)

Pseudo application of anaerobic metabolite:

The anaerobic metabolite is assumed to be applied directly to the soil by pseudo application. Hence, no “pathway” calculation was done in which the parent is applied. This is considered the only plausible but conservative way to account for the anaerobic formation (expressed by the maximum occurrence) and the aerobic degradation of the anaerobic metabolite. Applying the aerobic pathway for groundwater calculations may disregard the formation under anaerobic conditions.

Detailed application data used for simulation of  $PEC_{gw}$  for all compounds were compiled in Table CP 9.2.4.1- 12.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.4.1- 12: Application pattern used for PEC<sub>gw</sub> calculations

Individual crop	FOCUS crop used for interception	Application				Amount reaching soil per season application [g a.s./ha]
		Rate per season [g a.s. /ha]	Interval [days]	Plant interception [%]	BBCH stage	
Winter & spring cereals, GAP	-	2 × 150	14	-	30-69	-
Spring cereals 1, simulation	Spring cereals	2 × 150	14	2 × 80	30-69	2 × 30.0
Spring cereals 2, simulation <sup>2)</sup>	Spring cereals	2 × 22.68 <sup>1)</sup>	14	2 × 80	30-69	2 × 4.54 <sup>1)</sup>
Winter cereals 1, simulation	Winter cereals	2 × 150	14	2 × 80	30-69	2 × 30.0
Winter cereals 2, simulation <sup>2)</sup>	Winter cereals	2 × 22.68 <sup>1)</sup>	14	2 × 80	30-69	2 × 4.54 <sup>1)</sup>
Winter & spring cereals, GAP	-	2 × 125	14	-	30-61	-
Spring cereals 3, simulation	Spring cereals	2 × 125	14	2 × 80	30-61	2 × 25.0
Spring cereals 4, simulation <sup>2)</sup>	Spring cereals	2 × 18.90 <sup>1)</sup>	14	2 × 80	30-61	2 × 3.78 <sup>1)</sup>
Winter cereals 3, simulation	Winter cereals	2 × 125	14	2 × 80	30-61	2 × 25.0
Winter cereals 4, simulation <sup>2)</sup>	Winter cereals	2 × 18.90 <sup>1)</sup>	14	2 × 80	30-61	2 × 3.78 <sup>1)</sup>
Onions, GAP	-	2 × 125	10	-	15-47	-
Onions 1, simulation	Onions	2 × 125	10	2 × 10	15-47	2 × 112.5
Onions 2, simulation <sup>2)</sup>	Onions	2 × 18.90 <sup>1)</sup>	10	2 × 10	15-47	2 × 17.01 <sup>1)</sup>

<sup>1)</sup> Pseudo application is metabolite /ha

<sup>2)</sup> Pseudo application pattern for anaerobic metabolite HEC 5723 Carboxylic acid: parent rate – 1 d degradation, corrected for molar masses and maximum occurrence in anaerobic soil (= 100% metabolite rate)

For cereal and onion applications, absolute dates were derived for the simulation runs. All application dates are summarised in the table below.

Table CP 9.2.4.1- 13: Application dates and related information for fluoxastrobin as used for the simulation runs

Individual crop	Spring cereals	Winter cereals	Onions
Repeat Interval for App. Events	Every Year	Every Year	Every Year
Application Technique	Spray	Spray	Spray
Absolute / Relative to	Absolute	Absolute	Absolute
Scenario	1 <sup>st</sup> App. Date (Julian day)	1 <sup>st</sup> App. Date (Julian day)	1 <sup>st</sup> App. Date (Julian day)
	Chateaudun 10 Apr (100)	21 Apr (111)	29 May (149)

Substance specific and model related input parameters for FOCUS MACRO PEC<sub>gw</sub> calculations are summarised in Table CP 9.2.4.1- 14.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.4.1- 14: Compound input parameters for fluoxastrobin and its metabolites

Parameter	Unit	Fluoxastrobin (E+Z)	HEC 5725-E-des-chlorophenyl	HEC 5725-carboxylic acid	2-chlorophenol
<b>Common</b>					
Molar Mass	[g/mol]	458.8	348.3	417.8	128.56
Solubility	[mg/L]	2.292	9600	244 000	23 000
Vapour Pressure	[Pa]	5.63E-10	6.0E-05	7.00E-04	144
Freundlich Exponent		0.8584	0.9367	0.9043	0.8529
Plant Uptake Factor		0	0	0	0
Walker Exponent		0.49 <sup>1)</sup>	0.49 <sup>1)</sup>	0.49 <sup>1)</sup>	0.49 <sup>1)</sup>
DT <sub>50</sub>	[days]	38.89	56.7	17.0	23
Formation fraction		-	0.5145	-	1
<b>MACRO Parameters</b>					
K <sub>oc</sub>	[mL/g]	752.0	19.3	56.4	104.7
Q <sub>10</sub>		2.58 <sup>2)</sup>	2.58 <sup>2)</sup>	2.58 <sup>2)</sup>	2.58 <sup>2)</sup>
Canopy degradation half-life	[d]	10	10	10	10
Metabolite conversion factor (f <sub>convert</sub> ) <sup>3)</sup>		-	0.3906	-	0.2802

1) as proposed for MACRO 5.5.3

2) corresponding parameter in MACRO: tresp = 0.0948

3) metabolite formation in MACRO is based on molar masses M and formation fraction:  
f<sub>convert</sub> = M<sub>metab</sub> / M<sub>parent</sub> \* formation fraction

4) not available, as no formation fraction available, pseudo application used in MACRO

**Findings:** PEC<sub>GW</sub> were evaluated as the 80<sup>th</sup> percentile of the mean annual leachate concentration at 1 m soil depth. FOCUS MACRO PEC<sub>gw</sub> results for fluoxastrobin and its metabolites after application to winter and spring cereals and onions are given in the table below.

Table CP 9.2.4.1- 15: FOCUS MACRO PEC<sub>gw</sub> results of fluoxastrobin and its metabolites at Chateaudun

Scenario	Fluoxastrobin (E+Z)	HEC 5725-E-des-chlorophenyl	2-chlorophenol	HEC 5725-carboxylic acid <sup>1)</sup>
	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]	PEC <sub>gw</sub> [µg/L]
Spring cereals 2 × 150 g a.s./ha	<0.001	0.984	<0.001	<0.001
Winter cereals 2 × 150 g a.s./ha	<0.001	1.06	<0.001	<0.001
Spring cereals 2 × 125 g a.s./ha	<0.001	0.811	<0.001	<0.001
Winter cereals 2 × 125 g a.s./ha	<0.001	0.877	<0.001	<0.001
Onions 2 × 125 g a.s./ha	<0.001	3.81	<0.001	<0.001

1) Pseudo application pattern for the anaerobic metabolite HEC 5725-carboxylic acid (Scenario 1).

As described for scenario 1, 100 % of the potential pseudo application rate of anaerobic HEC 5725-carboxylic acid was applied, each year. All PEC<sub>gw</sub> values for all groundwater scenarios and application periods resulted already in concentrations < 0.001 µg/L, also without a division by 10. Therefore, a further simulation according Scenario 2, every 10 years, was not carried out anymore, as it is already covered with the first simulation.



**Conclusion:** There are no concerns for groundwater from the use of fluoxastrobin in accordance with the use pattern for the representative formulation.

The concentration of the metabolite HEC5725-E-des-chlorophenyl (M48) was predicted to reach groundwater at concentrations exceeding 0.1 µg/L. However, the relevance of this metabolite was assessed and the metabolite is non-relevant in groundwater (see Document N4).

#### CP 9.2.4.2 Additional field tests

No additional field studies were performed or required due to low PEC<sub>gw</sub> values calculated (see CP 9.2.4.1).

*This document is copyright protected.  
Any distribution, reproduction or publication requires the consent of Bayer AG (or its respective affiliate).  
Any use of the document or its content for regulatory or any other commercial purpose is prohibited and constitutes a violation of the underlying license agreement.*



**CP 9.2.5 Estimation of concentrations in surface water and sediment**

New calculations were performed, to reflect findings from new studies presented in the active substance dossier, section 7 “Fate and behaviour in the environment”. In addition these calculations consider the most recent guidance documents for exposure calculations. Calculations of predicted environmental concentrations are presented below.

**Predicted environmental concentrations in water (PEC<sub>sw</sub>) and sediment (PEC<sub>sed</sub>)**

**Endpoints for surface water (PEC<sub>sw</sub>) and sediment (PEC<sub>sed</sub>)**

For deriving the respective end points please refer to MCA Section 9, data point 7.2

**Table CP 9.2.5- 1: Key modelling input parameters for fluoxastrobin and its metabolites at Steps 1-2 level PEC calculations**

Parameter	Unit	Fluoxastrobin (E+Z)	HEC 5725 -E-des-chlorophenyl	HEC 5725 -carboxylic acid	2-chlorophenol
Molar Mass	g/mol	458.8	348.3	417.8	128.56
Water Solubility	mg/L	2.292	9600	244000	23000
Koc	mL/g	752	19.5	56.4	104.7
Degradation					
Soil	days	38.89	56.7	17.04	23
Total System	days	238.4	1000*	67.89	1000*
Water	days	238.4	1000	67.89	1000*
Sediment	days	1000*	1000*	67.89	1000*
Max Occurrence					
Water / Sediment	%	100	18.3	10.6	0.01
Soil	%	100	32.2	16.9	49.2

\* Default value used

**Table CP 9.2.5- 2: Additional modelling input parameters for fluoxastrobin at steps 3/4 level PEC calculations**

Parameter	Unit	Fluoxastrobin (E+Z)
General Parameters		
Molar Mass	g/mol	458.8
Water Solubility	mg/L	2.292
Vapour Pressure	Pa	5.6E-10
Plant Uptake Factor		0.0
Wash-Off Factor PRZM	l/cm	0.5
Wash-Off Factor MACRO	l/mm	0.05
Sorption		
Koc	mL/g	752
Freundlich Exponent		0.8584
Degradation		
Soil	days	38.89
Water	days	238.4
Sediment	days	1000
Water Exponent		0.7 (PRZM), 0.49 (MACRO)
Effect of Temperature		
Activation Energy	J/mol	65 400
Exponent	1/K	0.095
Q10		2.58





Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

**Report:** KCP 9.2.5/01 [redacted]; [redacted]; 2015; M-537907-01-1  
**Title:** Fluoxastrobin (FXA) and metabolites: PEC<sub>sw</sub>, sed FOCUS EUR - Use in cereals and onions in Europe  
**Report No.:** Ensa-15-0571  
**Document No.:** M-537907-01-1  
**Guideline(s):** not applicable  
**Guideline deviation(s):** not applicable  
**GLP/GEP:** no

**Materials and Methods:** Predicted environmental concentrations in surface water and sediment (PEC<sub>sw</sub> and PEC<sub>sed</sub>) of fluoxastrobin and its metabolites were calculated for the use in winter and spring cereals and onions in Europe. All relevant entry routes of a compound into surface water (combination of spray drift and runoff/erosion or drain flow) were considered in these calculations.

At FOCUS Step 2 the application period was set to March to May and the use in Northern and Southern Europe was considered. Details of the application pattern used in the Step 2 calculations are summarised in Table CP 9.2.5- 3.

**Table CP 9.2.5- 3: Application pattern used for PEC<sub>sw</sub> calculations at FOCUS Steps 1&2**

Crop	Rate [g a.s./ha]	Interval [days]	BBCH stage	FOCUS crop (crop group)	Season	Crop cover
Cereals, GAP	2 × 150	14	30-69	-	-	-
Cereals (winter), simulation 1	2 × 150	14	30-69	Winter cereals	Mar. - May	Intermediate crop cover 20 %
Cereals (spring), simulation 2	2 × 150	14	30-69	Spring cereals	Mar. - May	Intermediate crop cover 20 %
Cereals, GAP	2 × 125	14	30-61	-	-	-
Cereals (winter), simulation 1	2 × 125	14	30-61	Winter cereals	Mar. - May	Intermediate crop cover 20 %
Cereals (spring), simulation 2	2 × 125	14	30-61	Spring cereals	Mar. - May	Intermediate crop cover 20 %
Onions, GAP	2 × 125	10	15-47	-	-	-
Onions, simulation	2 × 125	10	15-47	Vegetables, bulb (arable crops)	Mar. - May	Minimal crop cover 10 %

In FOCUS Step 3 the application date for each scenario is determined by the Pesticide Application Timer (PAT), which is part of the FOCUS SW Scenarios. The user may only define an application time window. Absolute application dates for the crop simulation runs were estimated using a German regulatory tool AppDate 2<sup>2</sup>. Details of the parameters used in the Step 3 calculations are summarised in Table CP 9.2.5- 4.

<sup>2</sup> [redacted] 2015: Computer programme: "AppDate: Estimation of application dates based on crop development." (v.2.0b.).



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 4: Application dates of fluoxastrobin for the FOCUS Step 3 calculations

Parameter	Winter cereals	Spring cereals	Onions
PAT start date rel./absolute	Absolute	Absolute	Absolute
Appl. method (appl. type)	ground spray (CAM 2)	ground spray (CAM 2)	ground spray (CAM 2)
No of appl.	2	2	2
PAT window range	44	44	40
Appl. interval	14	14	10
Application Details	PAT Start Date	PAT Start Date	PAT Start Date
D1	20/04/02	27/05/01	21/04/01
D2	23/05/02	---	---
D3	02/07/02	28/04/01	18/04/01
D4	21/04/02	18/05/01	18/04/01
D5	15/03/02	09/04/01	---
D6, 1 <sup>st</sup>	02/03/02	---	05/05/01
D6, 2 <sup>nd</sup>	---	---	13/02/02
R1	20/04/02	---	15/04/01
R2	---	---	22/02/01
R3	10/04/02	---	22/02/01
R4	15/03/02	09/04/01	22/02/01

Compound input parameters for the Steps 1&2 simulation runs are summarised in Table CP 9.2.5- 1 and for the Steps 3&4 simulation runs in Table CP 9.2.5- 2.

**Findings: Steps 1&2:** The maximum PEC<sub>sw</sub> and PEC<sub>sed</sub> values for fluoxastrobin and its metabolites at Steps 1&2 are summarised in Table CP 9.2.5- 3.

This document is copyright protected. Any distribution, reproduction or publication of its content requires the consent of Bayer AG. Any use of the document for regulatory or commercial purposes is prohibited and constitutes a violation of the underlying license agreement.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 5: Maximum PEC<sub>sw</sub> and PEC<sub>sed</sub> values for fluoxastrobin and its metabolites at Steps 1&2

Use pattern	Scenario	Fluoxastrobin (E+Z)		HEC 5725 -E-des-chlorophenyl		HEC 5725 -carboxylic acid		2-chlorophenol	
		PEC <sub>sw</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]	PEC <sub>sw</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]	PEC <sub>sw</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]	PEC <sub>sw</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]
Cereals 2 × 150 g a.s./ha	Step 1	52.69	375.50	35.93	6.87	19.46	10.88	23.56	13.04
	Step 2								
	N-EU Multi	8.05	58.68	5.19	1.00	2.30	1.28	2.91	1.62
	S-EU Multi	14.66	108.41	10.10	0.95	4.48	2.51	5.63	3.14
	N-EU Single	4.54	33.09	2.86	0.55	1.41	0.79	1.78	0.98
S-EU Single	8.26	61.04	5.55	10.7	2.76	1.54	3.48	1.90	
Cereals 2 × 125 g a.s./ha	Step 1	43.91	312.92	29.94	5.73	16.22	9.08	19.63	10.95
	Step 2								
	N-EU Multi	6.71	48.90	4.33	0.83	1.91	1.07	2.43	1.35
	S-EU Multi	12.22	90.35	8.41	1.62	3.73	2.09	4.69	2.62
	N-EU Single	3.78	27.57	2.38	0.46	1.18	0.66	1.47	0.82
S-EU Single	6.88	50.88	4.62	0.89	2.20	1.29	2.99	1.11	
Onions 2 × 125 g a.s./ha	Step 1	43.91	312.92	29.94	5.73	16.22	9.08	19.63	10.95
	Step 2								
	N-EU Multi	7.39	54.08	4.84	0.93	2.14	1.15	2.84	1.58
	S-EU Multi	13.60	100.7	9.44	1.81	4.19	2.35	5.51	3.08
	N-EU Single	4.17	30.45	2.66	0.51	1.32	0.73	1.64	0.91
S-EU Single	7.66	56.69	5.18	1.00	2.58	1.44	3.18	1.77	

Step 3: The maximum PEC<sub>sw</sub>, PEC<sub>sed</sub> values and time weighted average concentrations at Day 7 of fluoxastrobin for relevant FOCUS Step 3 scenarios are given in the following tables.

Table CP 9.2.5- 6: Winter cereals: Maximum PEC<sub>sw</sub>, PEC<sub>sed</sub> and TWAC<sub>sw-7</sub> values for fluoxastrobin at Step 3

Use pattern	Fluoxastrobin (E+Z)					
	Cereals (winter), 2 × 150 g a.s./ha					
	Single application			Multiple applications		
FOCUS scenario	PEC <sub>sw</sub> [µg/L]	TWAC <sub>sw-7</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]	PEC <sub>sw</sub> [µg/L]	TWAC <sub>sw-7</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]
D1 (ditch)	0.048	0.825	3.289	1.044	0.874	5.479
D1 (stream)	0.864	0.110	1.304	0.795	0.272	2.590
D2 (ditch)	1.056	0.716	2.733	1.137	0.825	4.912
D2 (stream)	0.847	0.096	0.764	0.831	0.386	2.457
D3 (ditch)	0.952	0.499	0.692	0.834	0.200	0.865
D4 (pond)	0.033	0.030	0.254	0.042	0.040	0.443
D4 (stream)	0.731	0.0010	0.036	0.685	0.021	0.079
D5 (pond)	0.053	0.030	0.250	0.048	0.046	0.411
D5 (stream)	0.758	0.004	0.022	0.724	0.010	0.060
D6 (ditch)	0.948	0.147	0.553	0.834	0.353	1.021
R1 (pond)	0.076	0.072	0.696	0.203	0.193	1.683
R1 (stream)	0.625	0.071	0.532	1.663	0.207	1.522
R3 (stream)	0.883	0.113	1.268	1.337	0.182	1.429
R4 (stream)	0.807	0.218	0.836	1.724	0.483	1.931



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 7: Spring cereals: Maximum PEC<sub>sw</sub>, PEC<sub>sed</sub> and TWAC<sub>sw-7</sub> values for fluoxastrobin at Step 3

Use pattern	Fluoxastrobin (E+Z)					
	Cereals (spring), 2 × 150 g a.s./ha					
	Single application			Multiple applications		
FOCUS scenario	PEC <sub>sw</sub> [µg/L]	TWAC <sub>sw-7</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]	PEC <sub>sw</sub> [µg/L]	TWAC <sub>sw-7</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]
D1 (ditch)	1.010	0.826	3.916	1.403	1.204	7.135
D1 (stream)	0.841	0.137	1.595	0.728	0.220	3.184
D3 (ditch)	0.950	0.155	0.582	0.831	0.140	0.691
D4 (pond)	0.033	0.030	0.260	0.047	0.044	0.453
D4 (stream)	0.777	0.011	0.053	0.093	0.024	0.092
D5 (pond)	0.033	0.030	0.249	0.046	0.043	0.405
D5 (stream)	0.798	0.006	0.034	0.717	0.009	0.053
R4 (stream)	1.101	0.338	1.646	2.110	0.489	1.475

Table CP 9.2.5- 8: Winter cereals: Maximum PEC<sub>sw</sub>, PEC<sub>sed</sub> and TWAC<sub>sw-7</sub> values for fluoxastrobin at Step 3

Use pattern	Fluoxastrobin (E+Z)					
	Cereals (winter), 2 × 125 g a.s./ha					
	Single application			Multiple applications		
FOCUS scenario	PEC <sub>sw</sub> [µg/L]	TWAC <sub>sw-7</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]	PEC <sub>sw</sub> [µg/L]	TWAC <sub>sw-7</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]
D1 (ditch)	0.869	0.688	2.699	0.866	0.724	4.468
D1 (stream)	0.718	0.082	1.040	0.660	0.206	2.096
D2 (ditch)	0.870	0.586	2.196	0.936	0.676	3.951
D2 (stream)	0.706	0.071	0.601	0.683	0.313	1.946
D3 (ditch)	0.733	0.166	0.581	0.695	0.166	0.727
D4 (pond)	0.027	0.023	0.211	0.035	0.033	0.366
D4 (stream)	0.609	0.007	0.009	0.571	0.017	0.063
D5 (pond)	0.028	0.025	0.210	0.040	0.038	0.344
D5 (stream)	0.631	0.003	0.018	0.603	0.008	0.050
D6 (ditch)	0.790	0.120	0.463	0.695	0.294	0.859
R1 (pond)	0.062	0.059	0.580	0.167	0.158	1.404
R1 (stream)	0.521	0.058	0.449	1.355	0.169	1.281
R3 (stream)	0.736	0.092	1.060	1.090	0.149	1.196
R4 (stream)	0.662	0.180	0.702	1.410	0.397	1.620

This document is copyright protected. Any distribution or reproduction without the consent of Bayer AG (or its respective affiliates) for regulatory or constitutive purposes is prohibited and constitutes a violation of the underlying agreements.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 9: Spring cereals: Maximum PEC<sub>sw</sub>, PEC<sub>sed</sub> and TWAC<sub>sw-7</sub> values for fluoxastrobin at Step 3

Use pattern	Fluoxastrobin (E+Z)					
	Cereals (spring), 2 × 125 g a.s./ha					
	Single application			Multiple applications		
FOCUS scenario	PEC <sub>sw</sub> [µg/L]	TWAC <sub>sw-7</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]	PEC <sub>sw</sub> [µg/L]	TWAC <sub>sw-7</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]
D1 (ditch)	0.840	0.685	3.236	1.166	0.499	5.772
D1 (stream)	0.701	0.101	1.280	0.607	0.048	2.538
D3 (ditch)	0.792	0.129	0.488	0.693	0.117	0.581
D4 (pond)	0.027	0.025	0.216	0.039	0.036	0.374
D4 (stream)	0.647	0.008	0.044	0.078	0.010	0.076
D5 (pond)	0.028	0.025	0.209	0.038	0.036	0.339
D5 (stream)	0.665	0.005	0.039	0.597	0.007	0.044
R4 (stream)	0.900	0.277	1.578	1.780	0.401	0.065

Table CP 9.2.5- 10: Onions: Maximum PEC<sub>sw</sub>, PEC<sub>sed</sub> and TWAC<sub>sw-7</sub> values for fluoxastrobin at Step 3

Use pattern	Fluoxastrobin (E+Z)					
	Onions, 2 × 125 g a.s./ha					
	Single application			Multiple applications		
FOCUS scenario	PEC <sub>sw</sub> [µg/L]	TWAC <sub>sw-7</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]	PEC <sub>sw</sub> [µg/L]	TWAC <sub>sw-7</sub> [µg/L]	PEC <sub>sed</sub> [µg/kg]
D3 (ditch)	0.791	0.116	0.450	0.692	0.112	0.555
D4 (pond)	0.027	0.025	0.223	0.045	0.043	0.503
D4 (stream)	0.604	0.016	0.055	0.520	0.046	0.147
D6 (ditch)	0.788	0.059	0.278	0.693	0.160	0.630
D6 (ditch)	0.733	0.050	0.257	0.696	0.242	0.827
R1 (pond)	0.077	0.072	0.750	0.173	0.163	1.561
R1 (stream)	0.666	0.081	0.500	1.622	0.197	1.228
R2 (stream)	0.684	0.047	0.072	0.591	0.117	1.623
R3 (stream)	1.042	0.070	0.482	1.481	0.192	1.128
R4 (stream)	1.661	0.130	1.061	3.057	0.414	2.236

This document is copyright protected. Reproduction or its publication without the prior written consent of Bayer AG is prohibited. Any distribution, reproduction or use of this document without the prior written consent of Bayer AG is prohibited and constitutes a violation of the underlying patent law.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Step 4: The maximum PEC<sub>sw</sub> and PEC<sub>sed</sub> values and time weighted average concentrations at Day 7 of fluoxastrobin for relevant FOCUS Step 4 scenarios are given in the following tables.

Table CP 9.2.5- 11: Winter cereals: Maximum PEC<sub>sw</sub> values for fluoxastrobin at Step 4 after single and multiple applications

Buffer Width & Type		Scenario		Fluoxastrobin (E+Z)							
				Cereals (winter), 2 × 150 g a.s./ha							
				Single application				Multiple applications			
				PEC <sub>sw</sub> [µg/L] Drift Reduction				PEC <sub>sw</sub> [µg/L] Drift Reduction			
		0%	50%	75%	90%	0%	50%	75%	90%		
5m SD	D1 (ditch)	0.350	0.220	0.191	0.191	0.454	0.454	0.454	0.454		
	D1 (stream)	0.345	0.195	0.121	0.120	0.337	0.284	0.284	0.284		
	D2 (ditch)	0.454	0.454	0.454	0.454	0.951	0.951	0.951	0.951		
	D2 (stream)	0.336	0.289	0.289	0.289	0.597	0.597	0.597	0.597		
	D3 (ditch)	0.258	0.129	0.065	0.026	0.216	0.108	0.054	0.022		
	D4 (pond)	0.028	0.014	0.011	0.010	0.036	0.025	0.023	0.022		
	D4 (stream)	0.267	0.133	0.067	0.038	0.242	0.121	0.081	0.081		
	D5 (pond)	0.029	0.014	0.007	0.005	0.042	0.021	0.011	0.005		
	D5 (stream)	0.277	0.138	0.069	0.028	0.255	0.128	0.064	0.026		
	D6 (ditch)	0.257	0.129	0.064	0.026	0.216	0.108	0.054	0.033		
	R1 (pond)	0.074	0.037	0.064	0.062	0.199	0.182	0.182	0.178		
	R1 (stream)	0.571	0.571	0.571	0.571	1.663	1.663	1.663	1.663		
	R3 (stream)	0.809	0.809	0.809	0.809	1.337	1.337	1.337	1.337		
	R4 (stream)	0.807	0.807	0.807	0.807	1.724	1.724	1.724	1.724		
10m SD & RO	D1 (ditch)	0.228	0.191	0.191	0.191	0.454	0.454	0.454	0.454		
	D1 (stream)	0.205	0.125	0.120	0.120	0.284	0.284	0.284	0.284		
	D2 (ditch)	0.454	0.454	0.454	0.454	0.951	0.951	0.951	0.951		
	D2 (stream)	0.289	0.289	0.289	0.289	0.597	0.597	0.597	0.597		
	D3 (ditch)	0.137	0.068	0.034	0.014	0.112	0.056	0.028	0.011		
	D4 (pond)	0.020	0.011	0.010	0.010	0.027	0.024	0.022	0.021		
	D4 (stream)	0.147	0.071	0.038	0.033	0.125	0.081	0.081	0.081		
	D5 (pond)	0.021	0.010	0.005	0.002	0.030	0.015	0.008	0.004		
	D5 (stream)	0.147	0.077	0.037	0.015	0.133	0.066	0.033	0.021		
	D6 (ditch)	0.136	0.068	0.034	0.016	0.112	0.056	0.033	0.033		
	R1 (pond)	0.034	0.029	0.027	0.025	0.087	0.079	0.075	0.072		
	R1 (stream)	0.260	0.260	0.260	0.260	0.755	0.755	0.755	0.755		
	R3 (stream)	0.369	0.369	0.369	0.369	0.601	0.601	0.601	0.601		
	R4 (stream)	0.364	0.364	0.364	0.364	0.778	0.778	0.778	0.778		
20m SD & RO	D1 (ditch)	0.191	0.191	0.191	0.191	0.454	0.454	0.454	0.454		
	D1 (stream)	0.128	0.120	0.120	0.120	0.284	0.284	0.284	0.284		
	D2 (ditch)	0.454	0.454	0.454	0.454	0.951	0.951	0.951	0.951		
	D2 (stream)	0.289	0.289	0.289	0.289	0.597	0.597	0.597	0.597		
	D3 (ditch)	0.074	0.036	0.018	0.007	0.057	0.029	0.014	0.006		
	D4 (pond)	0.014	0.011	0.010	0.010	0.025	0.023	0.022	0.021		
	D4 (stream)	0.074	0.038	0.038	0.038	0.081	0.081	0.081	0.081		
	D5 (pond)	0.014	0.007	0.004	0.002	0.020	0.011	0.006	0.003		
	D5 (stream)	0.076	0.038	0.019	0.008	0.068	0.034	0.021	0.021		
	D6 (ditch)	0.071	0.035	0.018	0.016	0.057	0.033	0.033	0.033		
	R1 (pond)	0.019	0.015	0.014	0.013	0.046	0.041	0.038	0.036		
	R1 (stream)	0.136	0.136	0.136	0.136	0.395	0.395	0.395	0.395		
	R3 (stream)	0.194	0.194	0.194	0.194	0.314	0.314	0.314	0.314		
	R4 (stream)	0.190	0.190	0.190	0.190	0.406	0.406	0.406	0.406		

\* SD and RO denote spray drift and runoff buffer



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 12: Winter cereals: TWAC<sub>sw-7</sub> for fluoxastrobin at Step 4 after single and multiple applications

		Fluoxastrobin (E+Z)							
		Cereals (winter), 2 × 150 g a.s./ha							
		Single application				Multiple applications			
Buffer Width & Type	Scenario	TWAC <sub>sw-7</sub> [µg/L] Drift Reduction				TWAC <sub>sw-7</sub> [µg/L] Drift Reduction			
		0%	50%	75%	90%	0%	50%	75%	90%
5m SD	D1 (ditch)	0.283	0.183	0.176	0.176	0.435	0.435	0.435	0.435
	D1 (stream)	0.110	0.110	0.110	0.110	0.272	0.272	0.272	0.272
	D2 (ditch)	0.258	0.183	0.183	0.183	0.419	0.419	0.419	0.419
	D2 (stream)	0.096	0.096	0.096	0.096	0.233	0.233	0.233	0.233
	D3 (ditch)	0.054	0.027	0.013	0.005	0.052	0.026	0.008	0.005
	D4 (pond)	0.026	0.013	0.010	0.009	0.034	0.021	0.021	0.020
	D4 (stream)	0.010	0.010	0.010	0.010	0.021	0.021	0.021	0.021
	D5 (pond)	0.026	0.013	0.007	0.009	0.039	0.020	0.010	0.005
	D5 (stream)	0.001	0.001	0.000	0.000	0.004	0.002	0.002	0.002
	D6 (ditch)	0.040	0.020	0.010	0.004	0.091	0.045	0.022	0.009
	R1 (pond)	0.070	0.064	0.060	0.059	0.189	0.178	0.172	0.169
	R1 (stream)	0.071	0.071	0.071	0.071	0.207	0.207	0.207	0.207
	R3 (stream)	0.113	0.113	0.113	0.113	0.182	0.182	0.182	0.182
	R4 (stream)	0.218	0.218	0.218	0.218	0.483	0.483	0.483	0.483
10m SD & RO	D1 (ditch)	0.189	0.176	0.176	0.176	0.435	0.435	0.435	0.435
	D1 (stream)	0.110	0.110	0.110	0.110	0.272	0.272	0.272	0.272
	D2 (ditch)	0.183	0.183	0.183	0.183	0.419	0.419	0.419	0.419
	D2 (stream)	0.096	0.096	0.096	0.096	0.233	0.233	0.233	0.233
	D3 (ditch)	0.028	0.014	0.007	0.003	0.027	0.013	0.007	0.003
	D4 (pond)	0.010	0.010	0.009	0.009	0.025	0.022	0.021	0.020
	D4 (stream)	0.010	0.010	0.010	0.010	0.021	0.021	0.021	0.021
	D5 (pond)	0.019	0.009	0.005	0.002	0.028	0.014	0.008	0.004
	D5 (stream)	0.001	0.000	0.000	0.000	0.002	0.002	0.002	0.002
	D6 (ditch)	0.021	0.010	0.005	0.002	0.047	0.023	0.012	0.005
	R1 (pond)	0.032	0.028	0.025	0.024	0.083	0.075	0.071	0.068
	R1 (stream)	0.032	0.032	0.032	0.032	0.093	0.093	0.093	0.093
	R3 (stream)	0.052	0.052	0.052	0.052	0.081	0.081	0.081	0.081
	R4 (stream)	0.099	0.099	0.099	0.099	0.220	0.220	0.220	0.220
20m SD & RO	D1 (ditch)	0.176	0.176	0.176	0.176	0.435	0.435	0.435	0.435
	D1 (stream)	0.110	0.110	0.110	0.110	0.272	0.272	0.272	0.272
	D2 (ditch)	0.183	0.183	0.183	0.183	0.419	0.419	0.419	0.419
	D2 (stream)	0.096	0.096	0.096	0.096	0.233	0.233	0.233	0.233
	D3 (ditch)	0.045	0.007	0.004	0.001	0.014	0.007	0.003	0.001
	D4 (pond)	0.012	0.010	0.009	0.009	0.023	0.021	0.020	0.019
	D4 (stream)	0.010	0.010	0.010	0.010	0.021	0.021	0.021	0.021
	D5 (pond)	0.013	0.006	0.003	0.001	0.019	0.010	0.005	0.003
	D5 (stream)	0.000	0.000	0.000	0.000	0.002	0.002	0.002	0.002
	D6 (ditch)	0.011	0.005	0.003	0.002	0.024	0.012	0.006	0.004
	R1 (pond)	0.018	0.015	0.013	0.012	0.044	0.039	0.036	0.034
	R1 (stream)	0.017	0.017	0.017	0.017	0.049	0.049	0.049	0.049
	R3 (stream)	0.027	0.027	0.027	0.027	0.042	0.042	0.042	0.042
	R4 (stream)	0.052	0.052	0.052	0.052	0.115	0.115	0.115	0.115

\* SD and RO denote spray drift and runoff buffer



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 13: Winter cereals: Maximum PEC<sub>sed</sub> values for fluoxastrobin at Step 4 after single and multiple applications

Buffer Width & Type		Scenario		Fluoxastrobin (E+Z)								
				Cereals (winter), 2 × 150 g a.s./ha								
				Single application				Multiple applications				
				PEC <sub>sed</sub> [µg/kg] Drift Reduction				PEC <sub>sed</sub> [µg/kg] Drift Reduction				
				0%	50%	75%	90%	0%	50%	75%	90%	
5m SD	D1 (ditch)	2.370	2.321	2.296	2.281	4.739	4.609	4.543	4.503			
	D1 (stream)	1.300	1.298	1.298	1.298	2.579	2.575	2.574	2.573			
	D2 (ditch)	1.620	1.487	1.457	1.432	3.353	3.257	3.209	3.180			
	D2 (stream)	0.761	0.759	0.759	0.759	1.809	1.743	1.732	1.726			
	D3 (ditch)	0.197	0.101	0.052	0.021	0.239	0.123	0.064	0.027			
	D4 (pond)	0.228	0.145	0.103	0.079	0.399	0.219	0.198	0.160			
	D4 (stream)	0.035	0.035	0.034	0.034	0.045	0.074	0.073	0.073			
	D5 (pond)	0.218	0.115	0.062	0.029	0.358	0.194	0.111	0.066			
	D5 (stream)	0.008	0.004	0.002	0.002	0.022	0.012	0.007	0.007			
	D6 (ditch)	0.158	0.081	0.042	0.019	0.285	0.149	0.078	0.034			
	R1 (pond)	0.674	0.603	0.568	0.547	1.647	1.538	1.484	1.451			
	R1 (stream)	0.523	0.521	0.519	0.519	1.506	1.502	1.499	1.498			
	R3 (stream)	1.224	1.211	1.204	1.199	2.389	1.377	1.372	1.368			
	R4 (stream)	0.829	0.826	0.826	0.825	1.915	1.916	1.908	1.907			
10m SD & RO	D1 (ditch)	2.324	2.297	2.284	2.276	4.614	4.545	4.511	4.491			
	D1 (stream)	1.299	1.298	1.298	1.297	2.575	2.574	2.573	2.572			
	D2 (ditch)	1.490	1.459	1.443	1.434	3.261	3.211	3.186	3.171			
	D2 (stream)	0.760	0.759	0.759	0.758	1.744	1.733	1.727	1.723			
	D3 (ditch)	0.107	0.053	0.028	0.012	0.128	0.066	0.034	0.014			
	D4 (pond)	0.157	0.021	0.091	0.074	0.322	0.226	0.180	0.153			
	D4 (stream)	0.035	0.034	0.034	0.034	0.074	0.073	0.073	0.073			
	D5 (pond)	0.161	0.086	0.047	0.024	0.264	0.146	0.089	0.057			
	D5 (stream)	0.005	0.003	0.002	0.002	0.012	0.007	0.007	0.007			
	D6 (ditch)	0.086	0.045	0.024	0.011	0.154	0.081	0.043	0.023			
	R1 (pond)	0.436	0.282	0.255	0.238	0.777	0.694	0.652	0.627			
	R1 (stream)	0.183	0.182	0.180	0.181	0.508	0.506	0.504	0.504			
	R3 (stream)	0.352	0.340	0.341	0.338	0.458	0.452	0.449	0.447			
	R4 (stream)	0.366	0.365	0.364	0.364	0.805	0.802	0.801	0.800			
20m SD & RO	D1 (ditch)	2.298	2.284	2.277	2.273	4.547	4.512	4.495	4.484			
	D1 (stream)	1.298	1.298	1.297	1.297	2.574	2.573	2.573	2.572			
	D2 (ditch)	1.460	1.444	1.435	1.430	3.212	3.186	3.174	3.166			
	D2 (stream)	0.759	0.759	0.759	0.758	1.733	1.727	1.724	1.722			
	D3 (ditch)	0.057	0.029	0.015	0.006	0.067	0.035	0.018	0.007			
	D4 (pond)	0.142	0.101	0.082	0.070	0.256	0.194	0.164	0.146			
	D4 (stream)	0.034	0.034	0.034	0.034	0.073	0.073	0.073	0.073			
	D5 (pond)	0.111	0.069	0.034	0.019	0.183	0.106	0.071	0.050			
	D5 (stream)	0.003	0.002	0.002	0.002	0.007	0.007	0.007	0.007			
	D6 (ditch)	0.046	0.025	0.014	0.010	0.082	0.043	0.023	0.022			
	R1 (pond)	0.194	0.157	0.138	0.126	0.433	0.376	0.348	0.330			
	R1 (stream)	0.093	0.092	0.091	0.091	0.254	0.253	0.252	0.252			
	R3 (stream)	0.167	0.163	0.161	0.160	0.227	0.224	0.222	0.221			
	R4 (stream)	0.195	0.194	0.194	0.194	0.425	0.423	0.423	0.422			

\* SD and RO denote spray drift and runoff buffer





Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 14: Spring cereals: Maximum PEC<sub>sw</sub> values for fluoxastrobin at Step 4 after single and multiple applications

Buffer Width & Type		Scenario		Fluoxastrobin (E+Z)							
				Cereals (spring), 2 × 150 g a.s./ha							
				Single application				Multiple applications			
				PEC <sub>sw</sub> [µg/L] Drift Reduction				PEC <sub>sw</sub> [µg/L] Drift Reduction			
				0%	50%	75%	90%	0%	50%	75%	90%
5m SD	D1 (ditch)	0.311	0.235	0.235	0.235	0.523	0.523	0.523	0.523	0.523	0.523
	D1 (stream)	0.308	0.154	0.147	0.147	0.327	0.327	0.327	0.327	0.327	0.327
	D3 (ditch)	0.258	0.129	0.064	0.064	0.108	0.108	0.054	0.054	0.022	0.022
	D4 (pond)	0.028	0.014	0.012	0.012	0.040	0.028	0.026	0.026	0.024	0.024
	D4 (stream)	0.284	0.142	0.071	0.071	0.245	0.122	0.086	0.086	0.086	0.086
	D5 (pond)	0.029	0.015	0.007	0.007	0.039	0.020	0.010	0.010	0.005	0.005
	D5 (stream)	0.291	0.146	0.073	0.073	0.255	0.126	0.063	0.063	0.025	0.025
	R4 (stream)	1.101	1.101	1.101	1.101	2.177	2.177	2.177	2.177	2.177	2.177
10m SD & RO	D1 (ditch)	0.235	0.235	0.235	0.235	0.523	0.523	0.523	0.523	0.523	0.523
	D1 (stream)	0.164	0.147	0.147	0.147	0.327	0.327	0.327	0.327	0.327	0.327
	D3 (ditch)	0.137	0.068	0.034	0.034	0.112	0.056	0.028	0.028	0.011	0.011
	D4 (pond)	0.020	0.012	0.012	0.012	0.030	0.027	0.025	0.025	0.024	0.024
	D4 (stream)	0.151	0.075	0.043	0.043	0.127	0.086	0.086	0.086	0.086	0.086
	D5 (pond)	0.021	0.011	0.005	0.005	0.028	0.014	0.008	0.008	0.004	0.004
	D5 (stream)	0.155	0.077	0.039	0.039	0.131	0.066	0.033	0.033	0.019	0.019
	R4 (stream)	0.501	0.501	0.501	0.501	0.978	0.978	0.978	0.978	0.978	0.978
20m SD & RO	D1 (ditch)	0.235	0.235	0.235	0.235	0.523	0.523	0.523	0.523	0.523	0.523
	D1 (stream)	0.147	0.147	0.147	0.147	0.327	0.327	0.327	0.327	0.327	0.327
	D3 (ditch)	0.074	0.038	0.018	0.018	0.057	0.028	0.014	0.014	0.006	0.006
	D4 (pond)	0.014	0.012	0.011	0.011	0.028	0.025	0.024	0.024	0.024	0.024
	D4 (stream)	0.078	0.043	0.043	0.043	0.086	0.086	0.086	0.086	0.086	0.086
	D5 (pond)	0.014	0.007	0.004	0.004	0.019	0.010	0.005	0.005	0.003	0.003
	D5 (stream)	0.080	0.040	0.020	0.020	0.067	0.033	0.019	0.019	0.019	0.019
	R4 (stream)	0.262	0.262	0.262	0.262	0.510	0.510	0.510	0.510	0.510	0.510

\* SD and RO denote spray drift and runoff buffer

This document is copyrighted by Bayer AG for its use in the field of agricultural applications. Any distribution, reproduction or use of the document for any other commercial purpose is prohibited and constitutes a violation of the underlying license.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 15: Spring cereals: TWAC<sub>sw-7</sub> for fluoxastrobin at Step 4 after single and multiple applications

		Fluoxastrobin (E+Z)							
		Cereals (spring), 2 × 150 g a.s./ha							
		Single application				Multiple applications			
Buffer Width & Type	Scenario	TWAC <sub>sw-7</sub> [µg/L] Drift Reduction				TWAC <sub>sw-7</sub> [µg/L] Drift Reduction			
		0%	50%	75%	90%	0%	50%	75%	90%
5m SD	D1 (ditch)	0.256	0.219	0.219	0.219	0.512	0.512	0.512	0.512
	D1 (stream)	0.137	0.137	0.137	0.137	0.320	0.320	0.320	0.320
	D3 (ditch)	0.042	0.021	0.010	0.004	0.036	0.018	0.009	0.004
	D4 (pond)	0.026	0.013	0.011	0.010	0.037	0.026	0.024	0.023
	D4 (stream)	0.011	0.011	0.011	0.011	0.024	0.024	0.024	0.024
	D5 (pond)	0.026	0.013	0.007	0.003	0.037	0.024	0.010	0.004
	D5 (stream)	0.002	0.001	0.001	0.000	0.005	0.002	0.001	0.001
	R4 (stream)	0.338	0.338	0.338	0.338	0.479	0.476	0.474	0.473
10m SD & RO	D1 (ditch)	0.219	0.219	0.219	0.219	0.512	0.512	0.512	0.512
	D1 (stream)	0.137	0.137	0.137	0.137	0.320	0.320	0.320	0.320
	D3 (ditch)	0.022	0.011	0.006	0.002	0.019	0.009	0.005	0.002
	D4 (pond)	0.018	0.012	0.011	0.010	0.028	0.025	0.023	0.022
	D4 (stream)	0.011	0.011	0.011	0.011	0.024	0.024	0.024	0.024
	D5 (pond)	0.019	0.010	0.005	0.002	0.026	0.014	0.007	0.003
	D5 (stream)	0.001	0.001	0.000	0.000	0.002	0.001	0.001	0.001
	R4 (stream)	0.154	0.154	0.154	0.154	0.217	0.215	0.215	0.214
20m SD & RO	D1 (ditch)	0.219	0.219	0.219	0.219	0.512	0.512	0.512	0.512
	D1 (stream)	0.137	0.137	0.137	0.137	0.320	0.320	0.320	0.320
	D3 (ditch)	0.014	0.006	0.003	0.001	0.010	0.005	0.002	0.001
	D4 (pond)	0.011	0.011	0.010	0.010	0.026	0.024	0.023	0.022
	D4 (stream)	0.011	0.011	0.011	0.011	0.024	0.024	0.024	0.024
	D5 (pond)	0.013	0.006	0.003	0.002	0.018	0.009	0.005	0.003
	D5 (stream)	0.001	0.000	0.000	0.000	0.001	0.001	0.001	0.001
	R4 (stream)	0.081	0.081	0.081	0.081	0.113	0.113	0.112	0.112

\* SD and RO denote spray drift and runoff buffer

This document is copyrighted by its producer for regulatory and compliance purposes. Any distribution, reproduction or its use for any other commercial purpose is prohibited and constitutes a violation of the underlying license.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 16: Spring cereals: Maximum PEC<sub>sed</sub> values for fluoxastrobin at Step 4 after single and multiple applications

Buffer Width & Type		Scenario		Fluoxastrobin (E+Z)								
				Cereals (spring), 2 × 150 g a.s./ha								
				Single application				Multiple applications				
				PEC <sub>sed</sub> [µg/kg] Drift Reduction				PEC <sub>sed</sub> [µg/kg] Drift Reduction				
				0%	50%	75%	90%	0%	50%	75%	90%	
5m SD	D1 (ditch)	2.938	2.805	2.738	2.698	5.748	5.528	5.418	5.352			
	D1 (stream)	1.580	1.575	1.573	1.572	3.153	3.144	3.140	3.137			
	D3 (ditch)	0.165	0.084	0.043	0.018	0.190	0.098	0.050	0.021			
	D4 (pond)	0.234	0.152	0.110	0.086	0.410	0.281	0.218	0.181			
	D4 (stream)	0.040	0.039	0.039	0.038	0.086	0.084	0.083	0.082			
	D5 (pond)	0.217	0.115	0.063	0.031	0.352	0.190	0.106	0.060			
	D5 (stream)	0.013	0.007	0.004	0.002	0.020	0.010	0.006	0.006			
	R4 (stream)	1.631	1.627	1.624	1.623	2.445	2.437	2.433	2.430			
10m SD & RO	D1 (ditch)	2.813	2.742	2.707	2.686	5.536	5.422	5.365	5.331			
	D1 (stream)	1.576	1.573	1.572	1.571	3.144	3.140	3.138	3.136			
	D3 (ditch)	0.089	0.045	0.023	0.009	0.101	0.052	0.027	0.011			
	D4 (pond)	0.188	0.128	0.099	0.089	0.335	0.244	0.200	0.174			
	D4 (stream)	0.039	0.039	0.038	0.038	0.084	0.083	0.083	0.082			
	D5 (pond)	0.160	0.086	0.048	0.026	0.259	0.142	0.083	0.052			
	D5 (stream)	0.007	0.004	0.002	0.002	0.011	0.006	0.006	0.006			
	R4 (stream)	0.627	0.624	0.623	0.622	0.981	0.976	0.974	0.972			
20m SD & RO	D1 (ditch)	2.745	2.708	2.690	2.679	5.424	5.366	5.337	5.319			
	D1 (stream)	1.573	1.572	1.571	1.571	3.140	3.138	3.137	3.136			
	D3 (ditch)	0.047	0.024	0.012	0.005	0.052	0.027	0.014	0.006			
	D4 (pond)	0.150	0.108	0.089	0.078	0.272	0.214	0.185	0.168			
	D4 (stream)	0.039	0.038	0.038	0.038	0.083	0.083	0.082	0.082			
	D5 (pond)	0.111	0.061	0.035	0.021	0.179	0.101	0.065	0.044			
	D5 (stream)	0.004	0.002	0.002	0.002	0.006	0.006	0.006	0.006			
	R4 (stream)	0.324	0.323	0.322	0.322	0.511	0.509	0.508	0.507			

\* SD and RO denote spray drift and runoff buffer

This document is confidential or its publication or its content is prohibited and any distribution or use of the document for any other commercial purposes is a violation of the underlying license



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 17: Winter cereals: Maximum PEC<sub>sw</sub> values for fluoxastrobin at Step 4 after single and multiple applications

		Fluoxastrobin (E+Z)							
		Cereals (winter), 2 × 125 g a.s./ha							
		Single application				Multiple applications			
Buffer Width & Type	Scenario	PEC <sub>sw</sub> [µg/L] Drift Reduction				PEC <sub>sw</sub> [µg/L] Drift Reduction			
		0%	50%	75%	90%	0%	50%	75%	90%
5m SD	D1 (ditch)	0.287	0.178	0.141	0.141	0.348	0.348	0.348	0.348
	D1 (stream)	0.285	0.161	0.098	0.089	0.279	0.218	0.218	0.218
	D2 (ditch)	0.358	0.358	0.358	0.358	0.751	0.751	0.751	0.751
	D2 (stream)	0.274	0.229	0.229	0.229	0.474	0.474	0.474	0.474
	D3 (ditch)	0.215	0.107	0.054	0.022	0.180	0.098	0.045	0.018
	D4 (pond)	0.024	0.012	0.008	0.008	0.030	0.020	0.018	0.017
	D4 (stream)	0.223	0.111	0.056	0.031	0.201	0.101	0.065	0.065
	D5 (pond)	0.024	0.012	0.006	0.003	0.035	0.018	0.009	0.004
	D5 (stream)	0.231	0.115	0.058	0.023	0.213	0.106	0.055	0.021
	D6 (ditch)	0.214	0.107	0.054	0.021	0.180	0.090	0.045	0.026
	R1 (pond)	0.061	0.055	0.053	0.051	0.163	0.149	0.149	0.146
	R1 (stream)	0.466	0.466	0.466	0.466	1.355	1.355	1.355	1.355
	R3 (stream)	0.658	0.658	0.658	0.658	1.090	1.090	1.090	1.090
	R4 (stream)	0.662	0.662	0.662	0.662	1.410	1.410	1.410	1.410
10m SD & RO	D1 (ditch)	0.185	0.141	0.141	0.141	0.348	0.348	0.348	0.348
	D1 (stream)	0.168	0.102	0.089	0.089	0.218	0.218	0.218	0.218
	D2 (ditch)	0.358	0.358	0.358	0.358	0.751	0.751	0.751	0.751
	D2 (stream)	0.229	0.229	0.229	0.229	0.474	0.474	0.474	0.474
	D3 (ditch)	0.114	0.029	0.029	0.011	0.094	0.047	0.023	0.009
	D4 (pond)	0.009	0.009	0.008	0.008	0.022	0.019	0.018	0.017
	D4 (stream)	0.018	0.059	0.031	0.031	0.105	0.065	0.065	0.065
	D5 (pond)	0.017	0.009	0.004	0.002	0.025	0.013	0.007	0.003
	D5 (stream)	0.122	0.061	0.031	0.012	0.111	0.055	0.028	0.016
	D6 (ditch)	0.114	0.037	0.028	0.013	0.094	0.047	0.026	0.026
	R1 (pond)	0.028	0.024	0.022	0.021	0.072	0.065	0.062	0.059
	R1 (stream)	0.212	0.212	0.212	0.212	0.616	0.616	0.616	0.616
	R3 (stream)	0.300	0.300	0.300	0.300	0.490	0.490	0.490	0.490
	R4 (stream)	0.299	0.299	0.299	0.299	0.636	0.636	0.636	0.636
20m SD & RO	D1 (ditch)	0.141	0.141	0.141	0.141	0.348	0.348	0.348	0.348
	D1 (stream)	0.105	0.089	0.089	0.089	0.218	0.218	0.218	0.218
	D2 (ditch)	0.358	0.358	0.358	0.358	0.751	0.751	0.751	0.751
	D2 (stream)	0.229	0.229	0.229	0.229	0.474	0.474	0.474	0.474
	D3 (ditch)	0.059	0.030	0.015	0.006	0.048	0.024	0.012	0.005
	D4 (pond)	0.011	0.008	0.008	0.008	0.020	0.018	0.017	0.017
	D4 (stream)	0.061	0.031	0.031	0.031	0.065	0.065	0.065	0.065
	D5 (pond)	0.012	0.006	0.003	0.001	0.017	0.009	0.005	0.002
	D5 (stream)	0.064	0.032	0.016	0.006	0.056	0.028	0.016	0.016
	D6 (ditch)	0.059	0.030	0.015	0.013	0.048	0.026	0.026	0.026
	R1 (pond)	0.015	0.013	0.011	0.011	0.038	0.034	0.031	0.030
	R1 (stream)	0.111	0.111	0.111	0.111	0.322	0.322	0.322	0.322
	R3 (stream)	0.158	0.158	0.158	0.158	0.256	0.256	0.256	0.256
	R4 (stream)	0.156	0.156	0.156	0.156	0.332	0.332	0.332	0.332

\* SD and RO denote spray drift and runoff buffer



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 18: Winter cereals: TWAC<sub>sw-7</sub> for fluoxastrobin at Step 4 after single and multiple applications

Buffer Width & Type		Scenario		Fluoxastrobin (E+Z)							
				Cereals (winter), 2 × 125 g a.s./ha							
				Single application				Multiple applications			
				TWAC <sub>sw-7</sub> [µg/L] Drift Reduction				TWAC <sub>sw-7</sub> [µg/L] Drift Reduction			
				0%	50%	75%	90%	0%	50%	75%	90%
5m SD	D1 (ditch)	0.231	0.148	0.131	0.131	0.329	0.329	0.329	0.329	0.329	0.329
	D1 (stream)	0.082	0.082	0.082	0.082	0.206	0.206	0.206	0.206	0.206	0.206
	D2 (ditch)	0.205	0.135	0.135	0.135	0.330	0.330	0.330	0.330	0.330	0.330
	D2 (stream)	0.068	0.068	0.068	0.068	0.181	0.181	0.181	0.181	0.181	0.181
	D3 (ditch)	0.045	0.022	0.011	0.004	0.043	0.022	0.011	0.004	0.043	0.022
	D4 (pond)	0.021	0.011	0.008	0.007	0.028	0.016	0.017	0.016	0.028	0.016
	D4 (stream)	0.007	0.007	0.007	0.007	0.016	0.016	0.016	0.016	0.016	0.016
	D5 (pond)	0.022	0.011	0.006	0.002	0.033	0.017	0.009	0.004	0.033	0.017
	D5 (stream)	0.001	0.001	0.000	0.000	0.003	0.001	0.001	0.001	0.003	0.001
	D6 (ditch)	0.033	0.016	0.008	0.003	0.075	0.038	0.019	0.007	0.075	0.038
	R1 (pond)	0.058	0.052	0.050	0.048	0.155	0.146	0.141	0.138	0.155	0.146
	R1 (stream)	0.058	0.058	0.058	0.058	0.169	0.169	0.169	0.169	0.169	0.169
	R3 (stream)	0.092	0.092	0.092	0.092	0.149	0.149	0.149	0.149	0.149	0.149
	R4 (stream)	0.180	0.180	0.180	0.180	0.397	0.397	0.397	0.397	0.397	0.397
10m SD & RO	D1 (ditch)	0.153	0.131	0.131	0.131	0.329	0.329	0.329	0.329	0.329	0.329
	D1 (stream)	0.082	0.082	0.082	0.082	0.206	0.206	0.206	0.206	0.206	0.206
	D2 (ditch)	0.139	0.135	0.135	0.135	0.330	0.330	0.330	0.330	0.330	0.330
	D2 (stream)	0.068	0.068	0.068	0.068	0.181	0.181	0.181	0.181	0.181	0.181
	D3 (ditch)	0.024	0.022	0.006	0.002	0.022	0.011	0.006	0.002	0.022	0.011
	D4 (pond)	0.007	0.008	0.007	0.007	0.029	0.017	0.016	0.015	0.029	0.017
	D4 (stream)	0.007	0.007	0.007	0.007	0.016	0.016	0.016	0.016	0.016	0.016
	D5 (pond)	0.016	0.008	0.004	0.002	0.023	0.012	0.006	0.003	0.023	0.012
	D5 (stream)	0.001	0.000	0.000	0.000	0.002	0.001	0.001	0.001	0.002	0.001
	D6 (ditch)	0.017	0.009	0.004	0.002	0.039	0.019	0.010	0.004	0.039	0.019
	R1 (pond)	0.027	0.023	0.021	0.020	0.068	0.061	0.058	0.056	0.068	0.061
	R1 (stream)	0.026	0.026	0.026	0.026	0.076	0.076	0.076	0.076	0.076	0.076
	R3 (stream)	0.042	0.042	0.042	0.042	0.067	0.066	0.066	0.066	0.067	0.066
	R4 (stream)	0.082	0.082	0.082	0.082	0.181	0.181	0.181	0.181	0.181	0.181
20m SD & RO	D1 (ditch)	0.131	0.131	0.131	0.131	0.329	0.329	0.329	0.329	0.329	0.329
	D1 (stream)	0.082	0.082	0.082	0.082	0.206	0.206	0.206	0.206	0.206	0.206
	D2 (ditch)	0.135	0.135	0.135	0.135	0.330	0.330	0.330	0.330	0.330	0.330
	D2 (stream)	0.068	0.068	0.068	0.068	0.181	0.181	0.181	0.181	0.181	0.181
	D3 (ditch)	0.042	0.006	0.003	0.001	0.011	0.006	0.003	0.001	0.011	0.006
	D4 (pond)	0.010	0.008	0.007	0.007	0.018	0.016	0.016	0.015	0.018	0.016
	D4 (stream)	0.007	0.007	0.007	0.007	0.016	0.016	0.016	0.016	0.016	0.016
	D5 (pond)	0.010	0.005	0.003	0.001	0.015	0.008	0.004	0.002	0.015	0.008
	D5 (stream)	0.000	0.000	0.000	0.000	0.001	0.001	0.001	0.001	0.001	0.001
	D6 (ditch)	0.009	0.005	0.002	0.001	0.020	0.010	0.005	0.003	0.020	0.010
	R1 (pond)	0.014	0.012	0.011	0.010	0.036	0.032	0.030	0.028	0.036	0.032
	R1 (stream)	0.014	0.014	0.014	0.014	0.040	0.040	0.040	0.040	0.040	0.040
	R3 (stream)	0.022	0.022	0.022	0.022	0.035	0.035	0.035	0.035	0.035	0.035
	R4 (stream)	0.043	0.043	0.043	0.043	0.095	0.095	0.095	0.095	0.095	0.095

\* SD and RO denote spray drift and runoff buffer



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 19: Winter cereals: Maximum PEC<sub>sed</sub> values for fluoxastrobin at Step 4 after single and multiple applications

Buffer Width & Type		Scenario		Fluoxastrobin (E+Z)							
				Cereals (winter), 2 × 125 g a.s./ha							
				Single application				Multiple applications			
PEC <sub>sed</sub> [µg/kg] Drift Reduction				PEC <sub>sed</sub> [µg/kg] Drift Reduction							
0%	50%	75%	90%	0%	50%	75%	90%				
5m SD	D1 (ditch)	1.895	1.853	1.832	1.819	3.847	3.734	3.682	3.648		
	D1 (stream)	1.037	1.036	1.035	1.035	2.086	2.083	2.082	2.081		
	D2 (ditch)	1.257	1.182	1.157	1.142	2.638	2.557	2.516	2.492		
	D2 (stream)	0.598	0.597	0.597	0.596	1.400	1.358	1.349	1.343		
	D3 (ditch)	0.165	0.085	0.043	0.018	0.201	0.104	0.058	0.022		
	D4 (pond)	0.189	0.119	0.084	0.063	0.330	0.216	0.160	0.128		
	D4 (stream)	0.028	0.027	0.027	0.027	0.060	0.059	0.058	0.058		
	D5 (pond)	0.183	0.097	0.052	0.024	0.300	0.162	0.090	0.051		
	D5 (stream)	0.007	0.004	0.002	0.002	0.018	0.010	0.009	0.005		
	D6 (ditch)	0.132	0.068	0.035	0.015	0.240	0.125	0.065	0.028		
	R1 (pond)	0.563	0.503	0.473	0.455	1.374	1.282	1.236	1.209		
	R1 (stream)	0.441	0.439	0.438	0.435	1.268	1.265	1.263	1.262		
	R3 (stream)	1.023	1.011	1.005	1.002	2.162	1.152	1.147	1.145		
	R4 (stream)	0.696	0.694	0.693	0.692	1.606	1.600	1.600	1.599		
10m SD & RO	D1 (ditch)	1.855	1.853	1.822	1.815	3.742	3.684	3.655	3.637		
	D1 (stream)	1.036	1.035	1.035	1.035	2.083	2.082	2.081	2.080		
	D2 (ditch)	1.185	1.158	1.145	1.137	2.560	2.518	2.497	2.484		
	D2 (stream)	0.597	0.597	0.597	0.596	1.359	1.349	1.344	1.341		
	D3 (ditch)	0.089	0.048	0.023	0.010	0.108	0.056	0.029	0.012		
	D4 (pond)	0.150	0.099	0.074	0.059	0.265	0.183	0.144	0.121		
	D4 (stream)	0.027	0.027	0.027	0.027	0.059	0.058	0.058	0.058		
	D5 (pond)	0.135	0.072	0.039	0.020	0.220	0.121	0.071	0.044		
	D5 (stream)	0.004	0.002	0.002	0.002	0.010	0.006	0.005	0.005		
	D6 (ditch)	0.072	0.027	0.020	0.009	0.130	0.068	0.036	0.018		
	R1 (pond)	0.581	0.235	0.212	0.198	0.648	0.578	0.543	0.522		
	R1 (stream)	0.153	0.152	0.150	0.151	0.423	0.421	0.420	0.420		
	R3 (stream)	0.292	0.280	0.282	0.280	0.380	0.375	0.372	0.371		
	R4 (stream)	0.306	0.305	0.304	0.304	0.671	0.669	0.668	0.667		
20m SD & RO	D1 (ditch)	1.834	1.822	1.816	1.813	3.685	3.655	3.641	3.632		
	D1 (stream)	1.035	1.035	1.035	1.035	2.082	2.081	2.081	2.080		
	D2 (ditch)	1.159	1.146	1.139	1.135	2.518	2.497	2.486	2.480		
	D2 (stream)	0.597	0.596	0.596	0.596	1.349	1.344	1.342	1.340		
	D3 (ditch)	0.048	0.024	0.012	0.005	0.056	0.029	0.015	0.006		
	D4 (pond)	0.117	0.082	0.066	0.056	0.209	0.156	0.131	0.116		
	D4 (stream)	0.027	0.027	0.027	0.027	0.058	0.058	0.058	0.057		
	D5 (pond)	0.093	0.059	0.028	0.016	0.152	0.086	0.055	0.038		
	D5 (stream)	0.002	0.002	0.002	0.002	0.006	0.005	0.005	0.005		
	D6 (ditch)	0.039	0.020	0.011	0.008	0.069	0.036	0.019	0.016		
	R1 (pond)	0.163	0.131	0.115	0.105	0.362	0.314	0.290	0.275		
	R1 (stream)	0.077	0.076	0.076	0.076	0.211	0.210	0.210	0.209		
	R3 (stream)	0.138	0.134	0.133	0.131	0.188	0.185	0.184	0.183		
	R4 (stream)	0.163	0.162	0.162	0.162	0.354	0.353	0.352	0.352		

\* SD and RO denote spray drift and runoff buffer



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 20: Spring cereals: Maximum PEC<sub>sw</sub> values for fluoxastrobin at Step 4 after single and multiple applications

		Fluoxastrobin (E+Z)							
		Cereals (spring), 2 × 125 g a.s./ha							
		Single application				Multiple applications			
Buffer Width & Type	Scenario	PEC <sub>sw</sub> [µg/L] Drift Reduction				PEC <sub>sw</sub> [µg/L] Drift Reduction			
		0%	50%	75%	90%	0%	50%	75%	90%
5m SD	D1 (ditch)	0.256	0.178	0.178	0.178	0.411	0.411	0.411	0.411
	D1 (stream)	0.257	0.129	0.112	0.112	0.257	0.257	0.257	0.257
	D3 (ditch)	0.215	0.107	0.054	0.034	0.090	0.090	0.045	0.018
	D4 (pond)	0.024	0.012	0.009	0.009	0.033	0.022	0.020	0.019
	D4 (stream)	0.236	0.118	0.059	0.034	0.204	0.102	0.069	0.069
	D5 (pond)	0.024	0.012	0.006	0.003	0.033	0.009	0.009	0.004
	D5 (stream)	0.243	0.121	0.061	0.024	0.211	0.105	0.053	0.021
	R4 (stream)	0.900	0.900	0.900	0.900	1.786	1.786	1.786	1.786
10m SD & RO	D1 (ditch)	0.178	0.178	0.178	0.178	0.411	0.411	0.411	0.411
	D1 (stream)	0.136	0.112	0.112	0.112	0.257	0.257	0.257	0.257
	D3 (ditch)	0.114	0.057	0.028	0.011	0.093	0.047	0.023	0.009
	D4 (pond)	0.017	0.010	0.009	0.009	0.024	0.021	0.020	0.019
	D4 (stream)	0.125	0.063	0.034	0.034	0.106	0.069	0.069	0.069
	D5 (pond)	0.017	0.009	0.005	0.002	0.023	0.015	0.006	0.003
	D5 (stream)	0.129	0.064	0.032	0.013	0.109	0.055	0.027	0.015
	R4 (stream)	0.409	0.409	0.409	0.409	0.802	0.802	0.802	0.802
20m SD & RO	D1 (ditch)	0.178	0.178	0.178	0.178	0.411	0.411	0.411	0.411
	D1 (stream)	0.112	0.112	0.112	0.112	0.257	0.257	0.257	0.257
	D3 (ditch)	0.059	0.034	0.015	0.006	0.047	0.024	0.012	0.005
	D4 (pond)	0.009	0.009	0.009	0.009	0.022	0.020	0.019	0.019
	D4 (stream)	0.065	0.034	0.034	0.034	0.069	0.069	0.069	0.069
	D5 (pond)	0.012	0.006	0.003	0.001	0.016	0.008	0.004	0.002
	D5 (stream)	0.067	0.033	0.017	0.007	0.056	0.028	0.015	0.015
	R4 (stream)	0.214	0.214	0.214	0.214	0.419	0.419	0.419	0.419

\* SD and RO denote spray drift and runoff buffer

This document is copyrighted by Bayer AG or its affiliates. All rights reserved. Any distribution, reproduction or use of the document for any other commercial purpose is prohibited and may constitute a violation of the underlying license.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 21: Spring cereals: TWAC<sub>sw-7</sub> for fluoxastrobin at Step 4 after single and multiple applications

		Fluoxastrobin (E+Z)							
		Cereals (spring), 2 × 125 g a.s./ha							
		Single application				Multiple applications			
Buffer Width & Type	Scenario	TWAC <sub>sw-7</sub> [µg/L] Drift Reduction				TWAC <sub>sw-7</sub> [µg/L] Drift Reduction			
		0%	50%	75%	90%	0%	50%	75%	90%
5m SD	D1 (ditch)	0.211	0.163	0.163	0.163	0.398	0.398	0.398	0.398
	D1 (stream)	0.101	0.101	0.101	0.101	0.248	0.248	0.248	0.248
	D3 (ditch)	0.035	0.017	0.009	0.008	0.030	0.015	0.008	0.003
	D4 (pond)	0.021	0.011	0.009	0.008	0.031	0.021	0.019	0.018
	D4 (stream)	0.008	0.008	0.008	0.008	0.019	0.019	0.019	0.019
	D5 (pond)	0.022	0.011	0.006	0.002	0.031	0.020	0.008	0.004
	D5 (stream)	0.002	0.001	0.000	0.000	0.005	0.001	0.001	0.001
	R4 (stream)	0.277	0.277	0.277	0.277	0.393	0.390	0.389	0.388
10m SD & RO	D1 (ditch)	0.163	0.163	0.163	0.163	0.398	0.398	0.398	0.398
	D1 (stream)	0.101	0.101	0.101	0.101	0.248	0.248	0.248	0.248
	D3 (ditch)	0.018	0.009	0.005	0.002	0.016	0.008	0.004	0.002
	D4 (pond)	0.015	0.009	0.008	0.008	0.022	0.019	0.018	0.018
	D4 (stream)	0.008	0.008	0.008	0.008	0.019	0.019	0.019	0.019
	D5 (pond)	0.016	0.008	0.004	0.002	0.022	0.015	0.006	0.003
	D5 (stream)	0.001	0.001	0.000	0.000	0.001	0.001	0.001	0.001
	R4 (stream)	0.126	0.126	0.126	0.126	0.178	0.177	0.176	0.176
20m SD & RO	D1 (ditch)	0.163	0.163	0.163	0.163	0.398	0.398	0.398	0.398
	D1 (stream)	0.101	0.101	0.101	0.101	0.248	0.248	0.248	0.248
	D3 (ditch)	0.018	0.009	0.002	0.001	0.008	0.004	0.002	0.001
	D4 (pond)	0.015	0.009	0.008	0.008	0.020	0.019	0.018	0.017
	D4 (stream)	0.008	0.008	0.008	0.008	0.019	0.019	0.019	0.019
	D5 (pond)	0.011	0.005	0.003	0.001	0.015	0.008	0.004	0.002
	D5 (stream)	0.001	0.000	0.000	0.000	0.001	0.001	0.001	0.001
	R4 (stream)	0.066	0.066	0.066	0.066	0.093	0.092	0.092	0.092

\* SD and RO denote spray drift and runoff buffer

This document is reproduced for regulatory purposes only. Any distribution, reproduction or its use without the consent of Bayer AG is prohibited and constitutes a violation of the underlying license.





Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 22: Spring cereals: Maximum PEC<sub>sed</sub> values for fluoxastrobin at Step 4 after single and multiple applications

		Fluoxastrobin (E+Z)							
		Cereals (spring), 2 × 125 g a.s./ha							
Buffer Width & Type	Scenario	Single application				Multiple applications			
		PEC <sub>sed</sub> [µg/kg] Drift Reduction				PEC <sub>sed</sub> [µg/kg] Drift Reduction			
		0%	50%	75%	90%	0%	50%	75%	90%
5m SD	D1 (ditch)	2.365	2.253	2.197	2.163	4.656	4.471	4.379	4.323
	D1 (stream)	1.266	1.262	1.260	1.259	2.536	2.530	2.527	2.525
	D3 (ditch)	0.138	0.070	0.036	0.018	0.160	0.082	0.042	0.018
	D4 (pond)	0.194	0.124	0.089	0.069	0.339	0.229	0.175	0.144
	D4 (stream)	0.032	0.031	0.031	0.030	0.068	0.063	0.066	0.065
	D5 (pond)	0.182	0.097	0.053	0.025	0.296	0.192	0.088	0.048
	D5 (stream)	0.011	0.006	0.003	0.002	0.016	0.009	0.005	0.005
	R4 (stream)	1.366	1.362	1.360	1.359	2.040	2.033	2.030	2.027
10m SD & RO	D1 (ditch)	2.260	2.200	2.170	2.152	4.479	4.382	4.334	4.305
	D1 (stream)	1.263	1.261	1.260	1.259	2.530	2.527	2.525	2.524
	D3 (ditch)	0.074	0.038	0.019	0.008	0.085	0.044	0.023	0.009
	D4 (pond)	0.155	0.104	0.079	0.067	0.275	0.197	0.160	0.138
	D4 (stream)	0.031	0.031	0.030	0.030	0.067	0.066	0.065	0.065
	D5 (pond)	0.134	0.072	0.040	0.021	0.217	0.115	0.067	0.040
	D5 (stream)	0.006	0.003	0.002	0.002	0.009	0.005	0.005	0.005
	R4 (stream)	0.521	0.519	0.518	0.517	0.814	0.810	0.808	0.807
20m SD & RO	D1 (ditch)	2.202	2.171	2.156	2.146	4.384	4.335	4.311	4.296
	D1 (stream)	1.261	1.260	1.259	1.259	2.527	2.525	2.524	2.524
	D3 (ditch)	0.039	0.019	0.010	0.004	0.045	0.023	0.012	0.005
	D4 (pond)	0.152	0.088	0.071	0.062	0.277	0.172	0.147	0.133
	D4 (stream)	0.031	0.030	0.030	0.030	0.066	0.065	0.065	0.065
	D5 (pond)	0.093	0.051	0.029	0.017	0.149	0.083	0.052	0.034
	D5 (stream)	0.003	0.002	0.002	0.002	0.005	0.005	0.005	0.005
	R4 (stream)	0.269	0.268	0.267	0.267	0.424	0.421	0.420	0.420

\* SD and RO denote spray drift and runoff buffer

This document is copyrighted by Bayer AG. Any distribution, reproduction or its use for promotional or other commercial purposes without the consent of Bayer AG is prohibited and constitutes a violation of the underlying license.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 23: Onions: Maximum PEC<sub>sw</sub> values for fluoxastrobin at Step 4 after single and multiple applications

		Fluoxastrobin (E+Z)							
		Onions, 2 × 125 g a.s./ha							
Buffer Width & Type	Scenario	Single application				Multiple applications			
		PEC <sub>sw</sub> [µg/L] Drift Reduction				PEC <sub>sw</sub> [µg/L] Drift Reduction			
		0%	50%	75%	90%	0%	50%	75%	90%
5m SD	D3 (ditch)	0.214	0.107	0.054	0.021	0.180	0.090	0.045	0.018
	D4 (pond)	0.024	0.016	0.015	0.015	0.045	0.042	0.041	0.041
	D4 (stream)	0.221	0.110	0.055	0.043	0.185	0.112	0.112	0.112
	D6 (ditch)	0.213	0.107	0.054	0.025	0.180	0.090	0.045	0.025
	D6 (ditch)	0.213	0.107	0.054	0.025	0.181	0.090	0.046	0.026
	R1 (pond)	0.075	0.068	0.064	0.062	0.169	0.151	0.151	0.147
	R1 (stream)	0.666	0.666	0.666	0.666	1.622	1.622	1.622	1.622
	R2 (stream)	0.263	0.263	0.263	0.263	0.487	0.487	0.487	0.487
	R3 (stream)	1.042	1.042	1.042	1.042	1.481	1.481	1.481	1.481
	R4 (stream)	1.661	1.661	1.661	1.661	3.057	3.057	3.057	3.057
10m SD & RO	D3 (ditch)	0.114	0.057	0.028	0.011	0.093	0.047	0.023	0.009
	D4 (pond)	0.017	0.015	0.015	0.015	0.043	0.042	0.041	0.041
	D4 (stream)	0.117	0.059	0.043	0.043	0.112	0.112	0.112	0.112
	D6 (ditch)	0.113	0.057	0.029	0.025	0.094	0.048	0.025	0.025
	D6 (ditch)	0.113	0.057	0.029	0.014	0.094	0.048	0.026	0.026
	R1 (pond)	0.034	0.029	0.027	0.026	0.065	0.067	0.063	0.060
	R1 (stream)	0.302	0.302	0.302	0.302	0.734	0.734	0.734	0.734
	R2 (stream)	0.133	0.117	0.117	0.117	0.217	0.217	0.217	0.217
	R3 (stream)	0.467	0.466	0.466	0.466	0.675	0.675	0.675	0.675
	R4 (stream)	0.753	0.753	0.753	0.753	1.387	1.387	1.387	1.387
20m SD & RO	D3 (ditch)	0.059	0.030	0.015	0.006	0.047	0.024	0.012	0.005
	D4 (pond)	0.015	0.015	0.015	0.014	0.042	0.041	0.041	0.041
	D4 (stream)	0.061	0.043	0.043	0.043	0.112	0.112	0.112	0.112
	D6 (ditch)	0.059	0.030	0.025	0.025	0.048	0.025	0.025	0.025
	D6 (ditch)	0.059	0.030	0.015	0.014	0.048	0.026	0.026	0.026
	R1 (pond)	0.019	0.016	0.014	0.013	0.040	0.035	0.032	0.030
	R1 (stream)	0.158	0.158	0.158	0.158	0.384	0.384	0.384	0.384
	R2 (stream)	0.069	0.061	0.061	0.061	0.113	0.113	0.113	0.113
	R3 (stream)	0.243	0.243	0.243	0.243	0.354	0.354	0.354	0.354
	R4 (stream)	0.394	0.394	0.394	0.394	0.726	0.726	0.726	0.726

\* SD and RO denote spray drift and runoff buffer

Any distribution of this document without the consent of Bayer AG is prohibited. Any use of the document for purposes other than those intended by Bayer AG is a violation of the underlying license.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 24: Onions: TWAC<sub>sw-7</sub> for fluoxastrobin at Step 4 after single and multiple applications

Buffer Width & Type		Scenario		Fluoxastrobin (E+Z)							
				Onions, 2 × 125 g a.s./ha							
				Single application				Multiple applications			
		TWAC <sub>sw-7</sub> [µg/L] Drift Reduction				TWAC <sub>sw-7</sub> [µg/L] Drift Reduction					
		0%	50%	75%	90%	0%	50%	75%	90%		
5m SD	D3 (ditch)	0.031	0.016	0.008	0.003	0.029	0.015	0.007	0.003		
	D4 (pond)	0.021	0.015	0.014	0.014	0.042	0.040	0.039	0.039		
	D4 (stream)	0.016	0.016	0.016	0.016	0.045	0.045	0.045	0.045		
	D6 (ditch)	0.017	0.009	0.007	0.007	0.041	0.021	0.010	0.007		
	D6 (ditch)	0.014	0.012	0.012	0.012	0.063	0.038	0.019	0.019		
	R1 (pond)	0.070	0.063	0.060	0.058	0.159	0.141	0.138	0.138		
	R1 (stream)	0.081	0.081	0.081	0.081	0.197	0.197	0.197	0.197		
	R2 (stream)	0.047	0.047	0.047	0.047	0.117	0.117	0.117	0.117		
	R3 (stream)	0.078	0.078	0.078	0.078	0.192	0.192	0.192	0.192		
	R4 (stream)	0.175	0.175	0.175	0.175	0.414	0.414	0.414	0.414		
10m SD & RO	D3 (ditch)	0.017	0.008	0.004	0.002	0.015	0.008	0.004	0.002		
	D4 (pond)	0.015	0.014	0.014	0.014	0.041	0.040	0.039	0.039		
	D4 (stream)	0.016	0.016	0.016	0.016	0.045	0.045	0.045	0.045		
	D6 (ditch)	0.009	0.007	0.007	0.007	0.022	0.015	0.007	0.007		
	D6 (ditch)	0.012	0.012	0.012	0.012	0.033	0.019	0.019	0.019		
	R1 (pond)	0.032	0.028	0.025	0.024	0.061	0.063	0.058	0.056		
	R1 (stream)	0.036	0.036	0.036	0.036	0.088	0.088	0.088	0.088		
	R2 (stream)	0.021	0.021	0.021	0.021	0.053	0.053	0.053	0.053		
	R3 (stream)	0.036	0.036	0.036	0.036	0.088	0.088	0.088	0.088		
	R4 (stream)	0.079	0.079	0.079	0.079	0.186	0.186	0.186	0.186		
20m SD & RO	D3 (ditch)	0.009	0.004	0.002	0.001	0.008	0.004	0.002	0.001		
	D4 (pond)	0.014	0.014	0.014	0.013	0.040	0.039	0.039	0.038		
	D4 (stream)	0.016	0.016	0.016	0.016	0.045	0.045	0.045	0.045		
	D6 (ditch)	0.007	0.007	0.007	0.007	0.011	0.007	0.007	0.007		
	D6 (ditch)	0.012	0.012	0.012	0.012	0.019	0.019	0.019	0.019		
	R1 (pond)	0.018	0.015	0.013	0.012	0.038	0.032	0.030	0.028		
	R1 (stream)	0.019	0.019	0.019	0.019	0.046	0.046	0.046	0.046		
	R2 (stream)	0.011	0.011	0.011	0.011	0.028	0.028	0.028	0.028		
	R3 (stream)	0.019	0.019	0.019	0.019	0.046	0.046	0.046	0.046		
	R4 (stream)	0.041	0.041	0.041	0.041	0.097	0.097	0.097	0.097		

\* SD and RO denote spray drift and runoff buffer

Any distribution, reproduction or use of this document without the consent of Bayer AG is prohibited. Any other use of the document is a violation of the underlying license agreement.



Document MCP: Section 9 Fate and behaviour in the environment  
FXA+PTZ EC 200 (100+100) G

Table CP 9.2.5- 25: Onions: Maximum PEC<sub>sed</sub> values for fluoxastrobin at Step 4 after single and multiple applications

		Fluoxastrobin (E+Z)							
		Onions, 2 × 125 g a.s./ha							
Buffer Width & Type	Scenario	Single application				Multiple applications			
		PEC <sub>sed</sub> [µg/kg] Drift Reduction				PEC <sub>sed</sub> [µg/kg] Drift Reduction			
		0%	50%	75%	90%	0%	50%	75%	90%
5m SD	D3 (ditch)	0.127	0.065	0.033	0.014	0.152	0.078	0.040	0.017
	D4 (pond)	0.207	0.157	0.132	0.117	0.480	0.405	0.367	0.344
	D4 (stream)	0.054	0.053	0.053	0.053	0.145	0.144	0.144	0.144
	D6 (ditch)	0.093	0.062	0.047	0.042	0.256	0.150	0.096	0.067
	D6 (ditch)	0.732	0.675	0.646	0.629	1.533	1.442	1.399	1.372
	R1 (pond)	0.528	0.526	0.525	0.524	1.217	1.214	1.212	1.211
	R1 (stream)	0.669	0.668	0.668	0.668	1.618	1.616	1.615	1.615
	R2 (stream)	0.466	0.461	0.458	0.457	1.095	1.086	1.081	1.078
	R3 (stream)	1.046	1.041	1.038	1.037	2.214	2.208	2.200	2.203
	R4 (stream)	0.127	0.065	0.033	0.014	0.152	0.078	0.040	0.017
10m SD & RO	D3 (ditch)	0.068	0.035	0.018	0.007	0.081	0.042	0.022	0.009
	D4 (pond)	0.179	0.143	0.125	0.114	0.436	0.383	0.356	0.340
	D4 (stream)	0.053	0.053	0.053	0.053	0.144	0.144	0.144	0.144
	D6 (ditch)	0.064	0.048	0.042	0.042	0.155	0.098	0.069	0.067
	D6 (ditch)	0.354	0.310	0.288	0.275	0.720	0.651	0.617	0.596
	R1 (pond)	0.198	0.197	0.196	0.196	0.434	0.442	0.441	0.441
	R1 (stream)	0.172	0.171	0.171	0.171	0.408	0.407	0.406	0.406
	R2 (stream)	0.203	0.200	0.199	0.198	0.457	0.452	0.449	0.448
	R3 (stream)	0.418	0.413	0.413	0.413	0.864	0.861	0.859	0.858
	R4 (stream)	0.068	0.035	0.018	0.007	0.081	0.042	0.022	0.009
20m SD & RO	D3 (ditch)	0.036	0.018	0.009	0.004	0.042	0.022	0.011	0.005
	D4 (pond)	0.155	0.131	0.119	0.112	0.400	0.365	0.347	0.336
	D4 (stream)	0.053	0.053	0.053	0.053	0.144	0.144	0.144	0.144
	D6 (ditch)	0.048	0.042	0.042	0.042	0.099	0.070	0.067	0.067
	D6 (ditch)	0.201	0.170	0.155	0.145	0.399	0.352	0.329	0.315
	R1 (pond)	0.102	0.101	0.100	0.100	0.227	0.226	0.225	0.225
	R1 (stream)	0.079	0.079	0.079	0.079	0.186	0.185	0.185	0.185
	R2 (stream)	0.107	0.106	0.105	0.104	0.239	0.236	0.235	0.234
	R3 (stream)	0.215	0.213	0.213	0.212	0.445	0.443	0.442	0.442
	R4 (stream)	0.036	0.018	0.009	0.004	0.042	0.022	0.011	0.005

\* SD and RO denote spray drift and runoff buffer

Any distribution, reproduction or use of this document without the consent of Bayer AG is prohibited. Any other commercial use of the document is a violation of the underlying license agreement.



### CP 9.3 Fate and behaviour in air

For information on the fate and behaviour in air please refer to MCA Section 7, data point 7.3.

#### CP 9.3.1 Route and rate of degradation in air and transport via air

For information on route and rate of degradation in air and transport via air please refer to MCA Section 7, data points 7.3.1 and 7.3.2.

Due to the low volatility and short half-life in air no PEC calculations are required.

#### CP 9.4 Estimation of concentrations for other routes of exposure

There are no other routes of exposure if the product is used according to good agricultural practice. Therefore no further estimations are considered necessary.

*This document is copyright protected. Any distribution, reproduction or publication requires the consent of Bayer AG (or its respective affiliates). Any use of the document or its content for regulatory or other commercial purpose is prohibited and constitutes a violation of the underlying license agreement.*