

***Purpureocillium lilacinum* 251**
Microbial pest control agent against plant parasitic nematodes

Dossier according to OECD dossier guidance for microbial agents and microbial pest control products August 2006

Summary documentation Tier 1

Annex ICM, Section 5

Point IIM 7: Fate and Behaviour Studies on the Microbial Pest Control Agent in the Environment

Date: January 2016

Author:

Name:

Company:

Street no.:

Location:

Phone:

Fax:

Email:



M-544179-01-2

Table of Contents

IIM 7	Fate and Behaviour Studies on the Microbial Pest Control Agent in the Environment.....	5
IIM 7.1	Sufficient information on the origin, properties, survival and residual metabolites of the microorganism to assess its fate and behaviour in the environment. Viability/population dynamics, persistence, multiplication and mobility.....	5
IIM 7.1.1	Persistence and mobility in soil.....	5
IIM 7.1.2	Water.....	16
IIM 7.1.3	Air.....	18
IIM 7.2	Other/special studies.....	18
References	19

This document is copyright protected. Any distribution, reproduction or publication requires the consent of Bayer AG (or its respective affiliate). Any use of the document or its content for regulatory or commercial purposes is prohibited and constitutes a violation of the underlying license agreement.

OWNERSHIP STATEMENT

This document, the data contained in it and copyright therein are owned by Bayer CropScience. No part of the document or any information contained therein may be disclosed to any third party without the prior written authorisation of Bayer CropScience.

Any distribution, reproduction or publication requires the consent of Bayer AG (or its respective affiliate). Any use of the document or its content for regulatory or any other commercial purpose is prohibited and constitutes a violation of the underlying license agreement.

Introduction

The company Bayer CropScience AG is submitting a dossier for the re-approval of the microorganism *Purpureocillium lilacinum* 251 as an active substance under regulation (EC) 1107/2009.

The Microbial Pest Control Agent *Paecilomyces lilacinus* strain 251 was included into Annex I of Directive 91/414/EEC on 01/08/2008 (Commission Directive 2008/44/EC) and then approved according to the Commission Implementing Regulation (EU) No 540/2011 of 25 May 2011, implementing Regulation (EC) No 1107/2009 of the European Parliament¹. *P. lilacinus* strain 251 was notified and defended by Prophya GmbH. The active ingredient has been evaluated in Belgium according to Uniform Principles. The representative formulated product for the initial evaluation was the experimental formulation PBP-01001-I, containing 1×10^9 spores/g. PBP-01001-I, is comparable to the commercial formulation BioAct WG, containing 1×10^{10} spores/g, and the only changes between both formulations were slight adjustments of the content of two co-formulants, without any impact on the performance or physical properties of the formulated product. The recommended rate in terms of spores per hectare remained exactly the same. The data on PBP-01001-I can therefore be extrapolated to the formulated product BioAct WG, a wettable granule formulation (WG), the representative formulation in the present application for the renewal.

In 2013 Bayer CropScience AG acquired Prophya Biologischer Pflanzenschutz GmbH, now named Bayer CropScience Biologics GmbH. Bayer CropScience AG is the notifier for the renewal of *P. lilacinus* strain 251 in the procedure of AIR 3.

The microorganism has been previously classified as *Paecilomyces lilacinus* until 18S rRNA gene, internal transcribed spacer (ITS) and partial translation elongation factor α (TEF) sequencing revealed that *P. lilacinus* is not related to *Paecilomyces*. The new genus name *Purpureocillium* has been proposed for *P. lilacinus* and the new species name was assigned: *Purpureocillium lilacinum*. Therefore the strain is now identified as *Purpureocillium lilacinum*. In this dossier *Paecilomyces lilacinus* 251 and *Purpureocillium lilacinum* 251 are used as synonyms: *Paecilomyces lilacinus* = *Purpureocillium lilacinum*.

It has to be taken into account that data on *Paecilomyces lilacinus* from the open literature stated before 2011 may not necessarily provide reliable information due to insufficient classification methods used in these studies, especially, if the strain identification is not provided and/or identification methods used were based solely on morphological characteristics. However, they may provide relevant information transferrable to *Purpureocillium lilacinum*.

Purpureocillium lilacinum 251 is a ubiquitous, saprobic filamentous fungus commonly isolated from soil, decaying vegetation, insects and nematodes. Strains of *P. lilacinum* are used in plant protection products due to their nematicide activity. The mode of action against plant pathogenic nematodes of *P. lilacinum* strain 251 is principally based upon parasitism of nematode eggs as well as the vermiform stages of the nematodes, leading eventually to their death. With regard to the results of toxicity and ecotoxicity studies of the active substance *P. lilacinum* strain 251, it can be concluded that *P. lilacinum* strain 251 shows no risk for exposed humans, animals and environment.

P. lilacinum 251 is intended to be used in plant protection products to control plant pathogenic nematodes. The representative use presented in this dossier comprises applications of the formulation BioAct WG in protected and non-protected vegetable crops to control root-knot nematode, *Meloidogyne* spp.

Here we submit data that were previously evaluated by RMS Belgium as well as new data and information based on literature searches and studies.

¹ OJEU L94/13 Commission Directive 2008/44/EC of 4 April 2008 amending Council Directive 91/414/EEC to include benthiavalicarb, boscalid, carvone, fluoxastrobin, *Paecilomyces lilacinus* and prothioconazole as active substances

IIM 7 Fate and Behaviour Studies on the Microbial Pest Control Agent in the Environment

IIM 7.1 Sufficient information on the origin, properties, survival and residual metabolites of the microorganism to assess its fate and behaviour in the environment.

Viability/population dynamics, persistence, multiplication and mobility

New data 2015

Strain 251 was first described as *Paecilomyces lilacinus*. *P. lilacinus* is a common soil saprophyte and several strains belonging to this genus are aggressive parasites of plant parasitic nematodes. Strain 251 of *P. lilacinus* is the most well studied of the nematophagous strains and is commercialized for biocontrol in several countries. A new species, *Purpureocillium lilacinum*, was proposed in 2011 (██████████ et al., 2011, M-534512-01-1) as a result of phylogenetic analyses of 18S rRNA gene, internal transcribed spacer (ITS) and translation elongation factor 1- α (TEF) sequences. The phylogenetic data showed that *P. lilacinus* is not related to *Paecilomyces* and a new genus *Purpureocillium* was proposed with the type species *P. lilacinum*. Please refer also to Section 7, Point xx.

To gain sufficient information on fate and behaviour of *P. lilacinum* in soil, water and air, a literature search was performed using STN database (██████████ 2015, M-542861-01-1). 14 databases were considered in this search: Agricola, BIOSIS, MEDLINE, CAB Abstracts, SCISEARCH and Chemical Abstracts, DRUGU, EMBASE, Esbiobase, IPA, Pascal, POSiTech, Toxcenter and FSTA. After full text assessment, 15 articles published in the last ten years were determined to be relevant and were included in the dossier.

Cited references (abstracts):

Report: KIIM 7.1/01 – ██████████, J., ██████████, J., ██████████, T., ██████████, S-B., ██████████, A.M., ██████████, N.L., ██████████, R.A. (2011), *Purpureocillium*, a new genus for the medically important *Paecilomyces lilacinus*, published report

FEMS Microbiology Letters 321: 141-149

Abstract: *Paecilomyces lilacinus* was described more than a century ago and is a commonly occurring fungus in soil. However, in the last decade this fungus has been increasingly found as the causal agent of infections in man and other vertebrates. Most cases of disease are described from patients with compromised immune systems or intraocular lens implants. In this study, we compared clinical isolates with strains isolated from soil, insects and nematodes using 18S rRNA gene, internal transcribed spacer (ITS) and partial translation elongation factor 1- α (TEF) sequences. Our data show that *P. lilacinus* is not related to *Paecilomyces*, represented by the well-known thermophilic and often pathogenic *Paecilomyces variotii*. The new genus name *Purpureocillium* is proposed for *P. lilacinus* and the new combination *Purpureocillium lilacinum* is made here. Furthermore, the examined *Purpureocillium lilacinum* isolated grouped in two clades based on ITS and partial TEF sequences. The ITS and TEF sequences of the *Purpureocillium lilacinum* isolates used for biocontrol of nematode pests are identical to those causing infections in (immunocompromised) humans. The use of high concentrations of *Purpureocillium lilacinum* spores for biocontrol poses a health risk in immunocompromised humans and more research is needed to determine the pathogenicity factors of *Purpureocillium lilacinum*.

Report: KIIM 7.1/02 – ██████████, A. (2015) Literature review on *Purpureocillium lilacinum* strain 251: Fate and behaviour in the environment
non published report

Abstract: This report summarizes the search and selection process of open peer-reviewed literature for *Purpureocillium lilacinum* strain 251.

IIM 7.1.1 Persistence and mobility in soil

EU-Dossier: Doc M-IIB, Point 7.1.1

Persistence of *P. lilacinus* in the agricultural soil environment, into which it is delivered, is desired to accomplish efficacy and also is to be anticipated, since the soil is the original habitat of this saprophytic fungus.

For *P. lilacinus* and related species following information regarding persistence and survival in the soil environment can be derived from literature:

- Persistence of entomopathogenic Hyphomycetes in soil varies considerably on strain level, with *P. fumosoroseus* conidia being substantially degraded after 6 months (██████████ & ██████████, 1985, M-489363-01-1).
- Following application to soil some authors observed a substantial decline of several log in the content of viable spores of *P. lilacinus* per gram of soil, or complete clearance from spores within months after application (██████████ & ██████████, 1991, M-477635-01-1) but residual spores from a last year's application were also found at levels of 10^2 to 10^3 CFU/g of soil for different types of formulation employed (██████████ et al., 1999, M-477445-01-1).
- After planting tomatoes in a field which had been treated with *P. lilacinus* as a biocontrol agent for parasitic nematodes a year ago, residual populations of *P. lilacinus* were shown to increase by ~1 log within the growing season, up to harvest of these tomatoes (██████████ et al., 1999, M-477445-01-1), which apparently provide a source of parasitic nematodes for the fungus.
- The species *P. lilacinus* is considered a rather competitive organism in agricultural soils, based on development and research for biocontrol agents, and is assumed to readily establish in the micro-flora of soils, which also is supported by its adaptability to a wide range of soil pH (██████████ & ██████████, 1996, M-477546-01-1; ██████████, 1986, M-477553-01-1; ██████████ et al., 1981, M-477590-01-1).
- Survival of *P. lilacinus* is assumed to be as saprophyte (██████████ et al., 1999, M-477445-01-1), and the competitiveness was found not to be dependent on soil texture or composition of soils (██████████ & ██████████, 1987, M-489365-01-1).
- The type of fungal carrier used in the formulated product has a major effect on survival and efficacy of *P. lilacinus* spores, with granules being the most supportive ones for efficacy, and pellets supporting population establishment in soil (██████████ et al. 1989, M-489356-01-1).
- However, *P. lilacinus* may still be subject to soil biostatic factors, inhibiting germination (██████████ et al., 1980, M-476550-01-1) and also evidence for the zoopathogenic *Paecilomyces* species *P. fumosoroseus* and *P. farinosus* (██████████ & ██████████, 1984, M-477420-01-1).

Persistence of strain 251 of *P. lilacinus* in the agricultural soil environment

In the scope of her thesis on *P. lilacinus* strain 251 ██████████ performed three field trials on persistence of this fungus in natural soils, one of which is presented in ██████████ (1998, M-477414-01-1), showing a graph of decline of spore counts/gram garden soil within the first month after application:

Initially high levels of strain 251 of *P. lilacinus* in the range of 10^6 colony forming units/mL garden soil gradually declined within 18 days to 2.5 CFU/mL and after ~ five months the population had diminished to zero (see Fig. 7.1.1-01; based on data from ██████████ 2000, M-490110-01-1). Data on trial design and results for all three trials are presented in Table 7.1.1-01.




Figure 7.1.1-01: Persistence of *P. lilacinus* in garden soils (██████████, 2000, M-490110-01-1). DAT – days after treatment.

Table 7.1.1-01: Summary of persistence trials performed with *P. lilacinus* strain 251 (commercial product Paecil, 2×10^9 CFU/g) at different locations in Australia (██████████, 2000, M-490110-01-1).

Location	Trial design			Findings	
	Soil type, area	Application	Sampling, processing	Time sampling	Mean CFU/mL soil
#1. Garden at Macquarie University, Sydney	Undisturbed garden soil, 2x2m	18 g on 4 m ² = 4.5 g/m ² applied as soil drench in early December (additional irrigation)	Mixed 10 sub-samples of 10-16 cm top layer, 2 mL soil/sample was diluted and plated on semi-selective potato dextrose agar (PDA) for counting involving colony forming units/mL soil	Before application	0
				after application	1.3×10^6
				4 DAT	1.1×10^6
				6 DAT	7×10^5
				8 DAT	2.5×10^4
				32 DAT	2.5×10^4
				78 DAT	10^4
#2. 'Fauna Park' at Macquarie University, Sydney	Not specified garden soil 4.5 x 3.5m	15 g on 1.5 m ² = 10 g/m ² applied as soil drench in mid-December	Mixed 20 sub-samples of 10-20 cm top layer, 2 mL soil/sample was diluted and plated on semi-selective PDA for counting involving colony forming units/mL soil	before application	0
				after application	2.4×10^8
				7 DAT	5×10^4
				14 DAT	1.25×10^5
				28 DAT	0
				56 DAT	2.5×10^3
				119 DAT	0
#3. Vineyard near Broke, NSW	Not specified vine as soil drench in budburst	Mixed 6 sub-samples per vine from the rhizosphere. Inoculation plated on PDA for CFU counting	Before application	0	
			7 weeks after treatment (21 days)	In all 6 samples at 5×10^4 to 5×10^5	
			19 weeks after treatment (133 d)	In 3 of 6 samples at 2.5×10^4 to 5×10^5	

The results from the Fauna Park (#2) basically were in line with the findings at the garden plot (#1), with 6 colonies recovered 2 months after treatment, but some discrepancy is seen after 28 and 53 days with 80 colonies found after 4 weeks but a low level after 7-8 weeks. This variation in CFU counts can be explained by uneven distribution of fungal spores among the sampled area and by the additive systemic fault due to the low numbers of colonies found on agar plates (██████████, 2000, M-490110-01-1).

Recovery of *P. lilacinus* in the vineyard (#3) was much higher after 4-5 months following treatment, and showed no sign of a decline. According to ██████████ (2000, M-490110-01-1) this is presumably due to supporting weather conditions with light rain at the time of application, persisting for several days thereafter. Thus, spores found better starting conditions here, maybe also due to the root-knot nematodes present in the vineyard, being potential hosts for *P. lilacinus*. The deviation from the results of the soil sampling studies can also partly result from the different sampling site in the vineyard trial, which was the rhizosphere, a micro-habitat offering optimum conditions for growth of numerous saprophytic micro-organisms, including *P. lilacinus*. ██████████ (2000, M-490110-01-1) also stresses that the employed traditional method of serial dilution plating on semi-selective agar does not provide differentiation among *P. lilacinus* isolates, i.e. the biocontrol strain and native strains. In fact genetic analysis of some obtained cultures with differences in gross morphology showed that those isolates were definitely not strain 251. Therefore, the actual survival of strain 251 may be overestimated in these trials.

New data 2015

For *P. lilacinum* and related species following information regarding persistence and survival in the soil environment can be derived from literature search covering last 10 years, 2005-2015:

- [REDACTED], S. 2006; M-534735-01-1 described biological control of *Meloidogyne incognita* in laboratory experiments in tomato. The authors provide data on the survival of *P. lilacinum* strain 251 (PL251) after soil application, indicating an average density decrease of *P. lilacinum* strain 251 of 55% during the test period. Neither the presence of glucose, if applied as formulated product, nor the spore concentration had any effect on the persistence of *P. lilacinum* strain 251 in soil. This confirmed previous studies of the authors showing a drastical decline of *P. lilacinum* strain 251 in soil 14-21 days after application ([REDACTED] et al., 2004, cited in [REDACTED] et al., 2006).
- [REDACTED] & [REDACTED], 2006, M-534361-01-1 investigated on the effects of plant species on the persistence of *P. lilacinum* 251 in soil. 12 plant species, growing in soil pre-treated with *P. lilacinum* strain 251, were analysed. According to the results the tested plants had no effect on the behaviour of *P. lilacinum* strain 251 in soil. Furthermore, they showed a strong decline of the population density during testing, indicating a low potential for persistence and consequently for adverse environmental impacts. It was concluded that multiple applications of *P. lilacinum* strain 251 are necessary to maintain a high density for sufficient, long-term biocontrol ([REDACTED] & [REDACTED], 2006, M-534361-01-1). Furthermore, it could be demonstrated that the host plant is not the primary factor affecting the persistence of the fungus.
- [REDACTED] et al. (2005a, M-534360-01-1) evaluated the potential of strain PL251 to establish and survive in the environment after broadcast field application of a commercial water dispersible granule formulation (BIOACT WG) (4kg/ha). Within the first 90 days past application the density of *P. lilacinum* strain 251 decreased by more than 90%. At harvest, the *P. lilacinum* could no longer be isolated from the rhizosphere soil. It could be demonstrated that the decline in the population density of PL251 was independent from the spatial distribution and the population dynamic of the nematode *Heterodera schachtii*.
- [REDACTED] et al. (2005b, M-535178-01-1) evaluated the effects of the environment on the persistence of *P. lilacinum* strain 251 in soil in tomato and concluded that the strain shows relatively low persistence. The authors presented results that proved low environmental exposure to *P. lilacinum*, based on low field recovery after application (with 4×10^{13} conidia/ha applied and 5×10^5 expected, the recovery ranged between 10-50%); low persistence in soil and no growth after 144 h at 36°C.
- Better survival of *P. lilacinum* strain 251 was reported by [REDACTED] et al. (2015, M-534528-01-1) after repeated application in an organic citrus grove in Florida. The treatment resulted in *P. lilacinus* levels higher than those occurring naturally. Up to 12 months following its first application in the trial, *P. lilacinus* remained significantly more abundant in the *P. lilacinus* strain 251 plots than in the control. The authors did not provide any explanation for such a good persistence of *P. lilacinum* in citrus grove when compared to other crops.
- [REDACTED] (2006, M-534359-01-1) concluded from their experiments on *P. lilacinus* 251 that persistence and consequently the biocontrol efficacy of PL251 was not, unlike other nematophagous fungi, linked to the presence of the target nematode nor the host plant and that rhizosphere competence is not a key factor for the biocontrol efficacy of PL251. Multiple applications did increase the persistence of the fungus in soil which was correlated with excellent control of root-knot nematodes (*Meloidogyne incognita*) under field conditions.

- et al. (2008, M-534367-01-1) investigated the survival of *P. lilacinus* strain 251 (PL251) and the effect of application rate, substrate type, as well as the presence of the nematode host on its dynamics after application to the soil under controlled conditions. A gradual CFU decline was observed after application. The decline was independent from the application rates as well as from the presence of nematodes (Fig. 7.1.1-02). However, the substrate type had a significant effect on *P. lilacinum* persistence in soil. Clay soils favored fungus survival in comparison to sandy soils but also addition of organic matter to sandy soils (Fig. 7.1.1-03).




Fig. 7.1.1-02. Persistence of *P. lilacinum* strain 251 in the presence or absence of tomato plants and nematode host. Values are given as the mean of two application rates for each sampling (n=6) (after et al., 2008, M-534367-01-1).




Fig. 7.1.1-03. Persistence of *P. lilacinum* strain 251 in different soil substrates (after et al., 2008, M-534367-01-1).

- The above results were confirmed in another study on *P. lilacinum* strain 251 persistence using dilution plating techniques as well as molecular biology methods (nested PCR) for detection of fungus spores in the soil. In addition the interaction between *P. lilacinum* strain 251 and plants, nematodes, mutualistic fungal endophytes, and mycorrhiza was investigated. The studies showed that the initial density of the fungal antagonist after application was significantly lower than predicted and the spatial distribution was very heterogeneous. Already in the first year the density of PL251 already decreased by more than 90% within 90 days after application. After 120 days the fungus was no longer detected in soil. The observed decline was independent from the initial spatial distribution and not altered by the population dynamics of the host nematode. Two years after application the fungus was detected at equal level on treated and untreated plots; and the density of *P. lilacinum* strain 251 was far below the background level of other filamentous fungi. Furthermore, no adverse effects on mutualistic fungal endophytes, mycorrhiza, fungal antagonists or entomopathogenic nematodes were observed demonstrating the absence of competition under field conditions (2009 M-534891-01-1).
- [redacted] et al. (2005, M-534724-01-1), presented data on establishment of *P. lilacinum* on roots and soil in a period of 2 years, in carnation and gerbera plants. They found that the colonization with *P. lilacinum* increased significantly when the soil was pre-treated with dazomet, most likely due to reduction of competitor fungi.
- [redacted] et al. (2012, M-534851-01-1) showed that root-colonization of *P. lilacinus* in carrot field infested with *M. incognita* and *Erynia carotovora* was very efficient. After seed treatment (4×10^6 CFU /g seeds) application 10^2 CFU/g soil, and roots were recovered. Additionally the authors reported that root colonization by *P. lilacinus* and *Pseudomonas putida* was higher when they were applied together in comparison to individual treatments. *Pseudomonas putida* and *P. lilacinus* co-existed without affecting root colonization by either, which indicates that there is no antagonism between these two biocontrol species.
- [redacted] et al. (2015, M-53421-01-1) monitored the population of another strain of *P. lilacinus* (strain PL1210) in a field experiment for 60 days and showed that after an initial slight decrease from 1.2×10^5 spores/g soil to 0.3×10^5 spores/g soil, the fungus reached 0.99×10^5 after 60 days, which proves it was capable of colonizing the rhizosphere under the tested conditions (Fig. 7.1.1-04)




Fig. 7.1.1-04 Real-time quantification of *P. lilacinus* PL1210 collected at 1, 30, and 60 days after inoculation. The number of PL1210 spores was determined by plotting CT values against the standard curve.

Summary of persistence and mobility in soil - Persistence

Paecilomyces lilacinus is a saprophytic fungus naturally occurring in soil. Thus, its viability is naturally adapted to soil compartments. However, numerous studies have demonstrated that the survival of different strains of *P. lilacinus* in soil after application is limited in time. [redacted]; 2000; M-490110-01-1 investigated the persistence of *P. lilacinus* strain 251. In two sites, in a garden area and in a Fauna Park, they performed a quantitative examination of the fungus applied as the formulated product (WG formulation). Before application they could not detect any *P. lilacinus* viable spores. The strain 251 was then applied onto the surface of each a defined area and soil was sampled at several intervals until the level of viable fungi fell under the detection limit. It could be observed that the

number of *P. lilacinus* colony forming units (CFU) declined constantly within the first 2-3 weeks to less than 2% of the initially applied spores: from 1.3×10^6 down to 2.5×10^4 CFU/mL of soil in the garden area and from 2.4×10^8 down to 1.3×10^5 CFU/mL of soil in the Fauna Park, respectively.

These findings were supported by earlier studies. [REDACTED]; 1989; M-489356-01-1 conducted laboratory experiments to examine the survival of a peruvian strain of *P. lilacinus*. When the spores were applied to soil and incubated at room temperature for up to 56 days the decline was first slow but after 14 days the authors determined a fast decline.

[REDACTED], R.K.; M-477445-01-1 examined the efficacy of various plant leaves (castor, eucalyptus and neem), and *P. lilacinus* against *Meloidogyne incognita*. At the end of experiment, recovery of *P. lilacinus* was generally below detectable level or between 1 to 3×10^5 CFU/kg soil in treatment where it was used in combination with castor leaves, respectively. These findings also indicate a strong decline of viable spores in soil after application.

Apart from the data presented by [REDACTED] the available information provided for the first registration was based mainly on strains different from *P. lilacinum* strain PL251. In the meantime new supplemental information is available from public literature reporting on hosts with *P. lilacinum* strain 251 but also literature reporting on other *Paecilomyces* species. Generally the available new data provides comprehensive information on the behaviour of *P. lilacinum* strain 251 in soil and thus, should be mainly taken into account for the overall assessment. Data on other strains should only be considered if strain identification is possible. The newly available information was summarized above and generally confirms what was already concluded in the course of the first registration.

P. lilacinum strain 251 was reported to show a drastic decline within 14-21 days after a single soil application ([REDACTED], S. 2006, M-534735-01-1). Survival of *P. lilacinum* strain 251 in soil was independent from presence of plants ([REDACTED] & [REDACTED], 2006, M-534361-01-1) or an established rhizosphere. Furthermore, it was shown in this publication that also presence of host nematodes did not influence the survival of spores as a decline in the density of *P. lilacinum* strain 251 was reported independent from the spatial distribution of the target nematode *Heterodera schachtii*. The results were also confirmed in field trials where *P. lilacinum* strain 251 was shown to decrease by more than 90% within the first 90 days post application ([REDACTED] et al. 2005a, M-534360-01-1). Therefore, the authors concluded that multiple applications of *P. lilacinum* strain 251 are necessary to establish long-term nematode control. These findings are in line with results from [REDACTED] (2006, M-534359-01-1), showing better control of root-knot nematodes (*Meloidogyne incognita*) in field if *P. lilacinum* strain 251 is applied multiple times. Longer survival may also be due to organic matter (OM) present in soil. This is confirmed by studies from [REDACTED] et al. (2009, M-534367-01-1) showing better survival of *P. lilacinum* strain 251 if the OM content of soils is high. The content of organic matter could be one of the explanations for the good persistence of *P. lilacinum* observed in the citrus groves under organic agriculture in Florida, however this would require more studies ([REDACTED] et al., 2015).

Taken the information above into account it can be concluded that *P. lilacinum* strain 251 favours soil with a high organic content due to its saprophytic nature, and a better survival is achieved in such soils.

[REDACTED], 2009 M-534891-01-1 analysed the interaction between *P. lilacinum* strain 251 and plants, nematodes, mutualistic fungal endophytes and mycorrhiza and observed no adverse effects during testing, demonstrating the lack of competition under field conditions. Moreover, *P. lilacinum* strain 251 is under competition of natural occurring fungi. [REDACTED] et al. (2005, M-534724-01-1) showed an increased growth of *P. lilacinum* strain 251 in soil pre-treated with a fungicidal active product (azomet). Studies performed with *Paecilomyces lilacinus* ([REDACTED] et al., 2012), showed better survival in soil if applied in combination with *Pseudomonas putida* which may indicate synergism but not competitiveness of the two microorganisms. But since a different strain was tested this cannot be concluded for *P. lilacinum* strain 251. Thus, even if *P. lilacinum* strain 251 is applied on favourable soils, no adverse environmental impact can be expected and furthermore, competitiveness and persistence can be excluded.

In conclusion, following application of *P. lilacinum* strain 251 to soil the number of viable cells or spores of *P. lilacinus* are expected to show a fast decline to very low percentages within a few weeks. However, depending on the prevailing environmental conditions of the relevant soil ecosystem, they may possibly approach a balance at a clearly lower population density compared to the initial concentration, in response to limiting abiotic and also counteracting biotic factors. On a long-term scale,

without further applications of *P. lilacinum* strain 251, this saprophytic fungus may diminish completely, indicating the need for more than a single application to achieve nematode control. Therefore, since *P. lilacinum* strain 251 is naturally occurring in soil. Neither an unlimited multiplication nor an accumulation is expected.

Summary of persistence and mobility in soil - Mobility

Dispersal of spores of *P. lilacinum* strain 251 under conditions of use is limited, since it is intended to be applied directly onto the soil surface and incorporated by drench, or drip irrigation. Therefore dispersal via drift or via aerosols is not anticipated. Exposure to natural UV light will to some extent restrict germination and survival of applied *P. lilacinum* 251 into other environmental compartments, based on the results by [REDACTED], et al. (2000; M489366-01-1) for strain 251 tested for its UV sensitivity among other *P. lilacinus* strains. Since *P. lilacinum* strain 251 is a fungus dependent on aerobic respiration as well as its natural food, plant-parasitic nematodes, it is dependent on the upper aerobic zone of the soil. In deeper soil layers no survival of viable cells is expected. Due to their hydrophobicity the spores are expected to adsorb to soil particles and not to reach to lower zones.

In conclusion, there is no risk for uncontrolled growth of *P. lilacinus* in deeper soil layers, since this widespread and naturally occurring saprophytic soil fungus is subject to competition and antagonism in its natural habitat. Infectivity of strain 251 is confined to plant-parasitic nematodes and it is unable to grow at 37°C. Therefore any dispersal of this fungus imposes no health or environmental risk.

Cited references (abstracts):

Report: KIIM 7.1.1/15 – [REDACTED], S., [REDACTED], R.A. (2006), Biological control of the root-knot nematode *Meloidogyne incognita* by *Paecilomyces lilacinus* strain 251 published report
Biol Control 38: 179-187

Abstract: The fungal biocontrol agent, *Paecilomyces lilacinus* strain 251 (PL251), was evaluated for its potential to control the root-knot nematode *Meloidogyne incognita* on tomato. In growth chamber experiments, a pre-planting soil treatment reduced root galling by 66%, number of egg masses by 74% and the final nematode population in the roots by 71% compared to the inoculated control. Significant dose-response relationships were established when conidia were applied to soil either with or without the glucose-based formulation. The effective concentration₅₀ (EC₅₀) values for the commercially formulated product ranged between 0.097 g and 0.08 g/500 cm³ soil, equivalent to an EC₅₀ of 1.29 × 10⁶ and 9.88 × 10⁵ colony forming units (CFU)/g soil for the parameters gall index and final population per root, respectively. For the number of egg masses per root the EC₅₀ was 0.007 g product or 2.64 × 10⁵ CFU/g soil. Similarly, EC₅₀ values for conidia applied without formulation were 0.068 g or 0.103 g/500 cm³ soil (EC₅₀ of 8.10 × 10⁵–1.40 × 10⁶ CFU/g soil) for gall index and final population per root. In contrast, the EC₅₀ was 0.096 g (EC₅₀ of 1.25 × 10⁶ CFU/g soil) for the number of egg masses per root. We demonstrated that a single pre-plant application at a concentration of 1 × 10⁶ CFU/g soil is needed for sufficient biocontrol of *M. incognita* by PL251.

Report: KIIM 7.1.1/16 – [REDACTED], C., [REDACTED], S. (2006), Effect of plant species on persistence of *Paecilomyces lilacinus* strain 251 in soil and on root colonization by the fungus, published report
Plant and Soil 283: 25-34

Abstract: The effect of 12 plant species on the persistence of *Paecilomyces lilacinus* strain 251 in soil was investigated. After incorporating formulated conidia into non-sterile soil followed by transplanting different test plants, the population dynamic of the fungus was determined over 100 days. At termination of the experiment, the fungal population in the planted soil was compared to the density of *P. lilacinus* in the rhizosphere and the percent increase or decrease was calculated for each crop. In addition, the potential of *P. lilacinus* strain 251 to colonize roots endophytically was investigated. Comparison of the slopes describing the population dynamics of the fungus showed no significant differences between soil without plants and soil from the root zone of the majority of the test plants. Bean was the only plant species consistently exerting a negative effect on the persistence of *P. lilacinus* strain 251 in the soil. For the first time, *P. lilacinus* strain 251 was

isolated in significant numbers from healthy root tissue of barley plants.

Report: KIIM 7.1.1/17 – [REDACTED], S., [REDACTED], C., [REDACTED], R.A (2005a), Risk assessment of biological control products., published report
Gezunde Pflanzen 57: 163-166

Abstract: The egg pathogenic fungus *Paecilomyces lilacinus* strain 251 (PL251) can be used in an integrated approach to control the sugar beet cyst nematode *Heterodera schachtii*. To evaluate the potential of PL251 to establish and persist in the environment after broadcast application, a field experiment was conducted using a commercial water dispersible granule formulation (BIOACT WG). The fungal antagonist was applied at a rate of 4 kg product per ha and incorporated into the soil prior to planting sugar beets. At day zero, 50 and 90 post application and at harvest soil samples were collected to determine the population density of PL251. It was found that the spatial distribution after application was quite heterogeneous and that the density of the fungal antagonist directly after application was much lower than expected. Within the first 90 days post application the density of PL251 decreased more than 90 percent. At harvest, the antagonist could no longer be isolated from the rhizosphere soil. It could be demonstrated that the decline in the population density of PL251 was independent from the spatial distribution and the population dynamic of *H. schachtii*. Due to the fact that the fungal antagonist was not able to persist long under field conditions, the potential for PL251 to pose a risk to the environment is likely to be low.

Report: KIIM 7.1.1/18 – [REDACTED], [REDACTED], C., [REDACTED], R.A (2005b), Risk assessment of biological nematicides, published report
IOBC/wprs Bulletin 28: 201-205

Abstract: The development of a biological control product faces many obstacles before the final goal, successful commercialization, is achieved. Although several biopesticides have already proven their ability to efficiently control pests and diseases without causing any adverse effects to the environment, there are still concerns about the fate of a microorganism after its release. However, besides the monitoring of a specific biocontrol agent in the environment, models to appropriately assess the operator or bystander exposure need to be developed or modified for microbial pesticides. The egg pathogenic fungus *Paecilomyces lilacinus* (strain 251), was chosen as a model organism to identify the parameters needed to predict its fate in the environment. To monitor the long term survival, a semi-selective medium was developed to enable monitoring of the population dynamics of *P. lilacinus* in the soil and rhizosphere. Monitoring was conducted with root knot nematodes (*Meloidogyne* spp.) as target and tomato as host plant. The population development of *P. lilacinus* was monitored depending on the application rate, formulation, temperature, method of application and the presence or absence of the target pest as well as the host plant. Initial results demonstrated that *P. lilacinus* was not able to multiply in soil or the rhizosphere of tomato plants and consequently showed a relatively low persistence...

Report: KIIM 7.1.1/19 – [REDACTED], R.; [REDACTED], F. E.; [REDACTED], L. W. (2015)
Modifying soil to enhance biological control of belowground dwelling insects in citrus groves under organic agriculture in Florida, published report
Biological Control 84: 53-66

Abstract: An emerging organic citrus industry in Florida could benefit greatly from effective, non-conventional methods to mitigate losses from pests and diseases. We studied part of a soil food web in an organic orchard to learn ways to conserve and enhance biological control of insect pests by native entomopathogenic nematodes (EPNs). We evaluated two OMRI (Organic Materials Review Institute) approved cultural practices: (i) a mulch of commercially pelleted chicken manure, (ii) a commercial formulation of *Purpureocillium lilacinus*, and (iii) an un-amended control. Several soil nutrients (i.e. nitrogen, phosphate, and potassium) were affected by the amendments, but initial equilibrium values (T0) were restored by the last sampling time (T12). The plant parasitic nematode *Tylenchulus semipenetrans* increased in both treatments compared to the untreated control at T3 ($P < 0.05$). The oomycete *Phytophthora nicotianae* increased in the *P. lilacinus* plots at T1, marginally at T12, but decreased at T6 and T9. *Steinernema diaprepesi*, *Heterorhabditis indica* and *Heterorhabditis zealandica* were the only EPNs regularly detected in the orchard. Mulch increased numbers of *H. zealandica* at T6 and T9 ($P < 0.05$) and free living nematodes at T12 ($P < 0.01$). The nematophagous fungus (NF) *P. lilacinus* persisted in plots where it was augmented ($P < 0.05$), reaching a maximum level at T3 that was 17.5-fold greater than that in controls. Numbers of *Paenibacillus* sp. were directly related to both those of *S. diaprepesi* and

Acrobeloides-group nematodes ($P < 0.01$), but inversely to the FLN counts ($P < 0.05$). The application of these two amendments did not produce strong changes in the EPN community but decreased the emergence from soil of adult *Diaprepes abbreviatus*, a root weevil pest. Thus, both amendments might contribute to citrus pest management under organic production.

Report: KIIM 7.1.1/20 – [REDACTED], S., 2006b Multitrophic interactions of *Paecilomyces lilacinus* strain 251 in the rhizosphere of host and non-host plants.

published report

IOBC/wprs Bulletin 29: 53-61

Abstract: The facultative egg pathogenic fungus *Paecilomyces lilacinus* is one of the most widely tested biocontrol agents for control of plant parasitic nematodes. The commercial strain 251 (PL251) is undergoing registration procedures in the EU and US and is commercially available as BIOACT® WG in several countries. To better understand the multitrophic interactions of PL251 in the rhizosphere, dose-response experiments were conducted to evaluate the relationship between the antagonist dose and biocontrol efficacy and fungal persistence. The importance of host- or non-host plants, nematodes, mutualistic fungal endophytes, and mycorrhiza for biological efficacy and unwanted side effects caused by the application of the biocontrol fungus was also investigated. It could be demonstrated that persistence and consequently the biocontrol efficacy of PL251 is not, unlike other nematophagous fungi, linked to the presence of the target nematode nor the host plant. Furthermore, some nematode host plants seem to provide unsuitable conditions in their rhizosphere resulting in rapid decline of fungal density and in some cases reduced efficacy of the antagonist. In contrast to other nematophagous fungi, rhizosphere competence is not a key factor for the biocontrol efficacy of PL251. Multiple applications did increase the persistence of the fungus in soil which was correlated with excellent control of root-knot nematodes under field conditions.

Report: KIIM 7.1.1/21 – [REDACTED], C., [REDACTED], A., [REDACTED], R.A., [REDACTED], S. (2008), Persistence of the nematophagous fungus *Paecilomyces lilacinus* strain 251 in soil under controlled conditions.

published report

Biocontrol Sci and Tech 18: 941-1056

Abstract: The persistence of the nematophagous fungus *Paecilomyces lilacinus* [REDACTED] strain 251 (PL251) and the effect of application rate, substrate type, as well as the presence of the nematode host on its dynamics after application to the soil were investigated under controlled conditions. In all experiments, increase of *P. lilacinus* colony forming units after application was not found. In contrast a gradual decline in fungal densities over time was observed. Application rate had no significant effect on the dynamics of the fungal population. Likewise, *P. lilacinus* density decline in soil was not significantly affected by the presence of the nematode host. Substrate type had a significant effect on *P. lilacinus* persistence in soil. The fungal agent persisted longer in silty loam and clay soil, with reduced persistence when sand was added to field soil. Conversely, when organic substrate was added to pure sand, persistence was significantly increased. Although persistence of fungal biocontrol agents in soil depends on various biotic and abiotic conditions, baseline data on persistence such as those reported in this study are helpful for biocontrol and environmental risk assessment and merit further study.

Report: KIIM 7.1.1/22 – [REDACTED], S. (2009), Understanding multitrophic interactions to facilitate successful biocontrol of plant parasitic nematodes with *Paecilomyces lilacinus* strain 251.

published report

IOBC/wprs Bull 43: 297-298

Abstract: The facultative egg-pathogenic fungus *Paecilomyces lilacinus* strain 251 (PL251) is one of the most widely tested biocontrol agents for control of plant-parasitic nematodes. Recently, PL251 was included as active substance in Annex I to the directive 91/414/EEC. In the USA, PL251 is registered as bio-nematicide under the trade name MELOCON® WG for use on a variety of crops. So far PL251 has demonstrated efficacy in reducing root-knot, cyst and free living plantparasitic nematodes on a range of crops. However, to better understand the multitrophic interactions of PL251 with host- or non-host plants, nematodes, mutualistic fungal endophytes, and mycorrhiza studies were conducted to determine their importance for biological efficacy. In none of the studies conducted, adverse effects on mutualistic fungal endophytes, mycorrhiza, fungal antagonists or entomopathogenic nematodes were observed. Conversely to other nematophagous fungi, rhizosphere competence seems not a key factor for the efficacy of PL251. However, studies

are underway to determine the eggmass colonisation by PL251 using realtime PCR assays which are able to detect 10 CFU per eggmass or less. Monitoring the persistence of PL251 under field conditions using dilution plating techniques and nested PCR revealed a rapid decline of the fungal density in soil over time. Although detection of PL251 in soil was still possible two years after application, the overall suppressiveness of egg pathogenic fungi towards cyst nematodes was not affected.

Report: KIIM 7.1.1/23 – [REDACTED], M., [REDACTED], P.P. (2005), Management of carnation and gerbera to control the root-knot nematode, *Meloidogyne incognita*, in commercial polyhouses. published report
Nematol medit 33: 157-162

Abstract: Studies on the management of carnation and gerbera to control root-knot nematode, *Meloidogyne incognita*, in commercial polyhouses using pre-plant (dazomet) and post-plant (chlorpyrifos, carbosulfan and carbofuran) chemicals in a comparison with various combinations of and with bio-agents (*Paecilomyces lilacinus*, *Pochonia chlamydosporia*) and neem cake were made. Pre-plant treatment of beds with dazomet followed by the application of neem cake (1 kg/m², 15 days later) along with *P. lilacinus* or *P. chlamydosporia* significantly reduced populations of *M. incognita* and the mortality of plants, and suppressed the nematode infection for nearly 2 years. The antagonistic fungi established themselves better in the beds treated with dazomet than in untreated beds. Chlorpyrifos and carbofuran (each applied twice in 6 months) significantly reduced nematode populations in roots and soil. However, there was a build-up of nematode populations in beds treated with these two chemicals after 1 year. On a long term basis, soil management with pre-plant treatment of dazomet followed by the application of oil cakes plus antagonistic fungi, was more effective against *M. incognita* than postplant treatment with carbofuran, carbosulfan and chlorpyrifos on carnation and gerbera grown in polyhouses.

Report: KIIM 7.1.1/24 – [REDACTED], D., [REDACTED], M. S., [REDACTED], R. M., [REDACTED], J., [REDACTED], K. (2012), Bio-management of *Meloidogyne incognita* and *Erwinia carotovora* in carrot (*Daucus carota* L.) using *Pseudomonas putida* and *Paecilomyces lilacinus*. published report
Nematol medit 40: 189-194

Abstract: An experiment was conducted to test the effects of two biocontrol agents, *Pseudomonas putida* and *Paecilomyces lilacinus*, on the control of *Meloidogyne incognita* and *Erwinia carotovora* in carrot (*Daucus carota*). *Pseudomonas putida* and *P. lilacinus* formulations were enriched in neem cake and evaluated under field conditions, individually and in combination, as seed treatment or as substrate treatment. Twenty grams of this formulation was used for the seed (one kg) treatment and five kg for the enrichment of neem cake (200 kg), which was applied to the beds at the rate of 20 g/m² as a substrate treatment before sowing. Seed treatment with both bio-agents and application of neem cake enriched with *P. putida* and *P. lilacinus* proved to be the best of all the treatments, leading to a reduction in the *M. incognita* (J2) population (in roots by 69 percent and soil by 47.6 percent) and *E. carotovora* by 66 percent, with a significant increase (27.8 percent) in the yield of carrot. *Pseudomonas putida* and *P. lilacinus* co-existed without affecting root colonisation by either.

Report: KIIM 7.1.1/25, [REDACTED], Z., [REDACTED], Y., [REDACTED], W., [REDACTED], Y. 2015 Root colonization and effect of biocontrol fungus *Paecilomyces lilacinus* on composition of ammonia-oxidizing bacteria, ammonia-oxidizing archaea and fungal populations of tomato rhizosphere. published report
Biol Fertil Soils 51:343-351

Abstract: This study investigated the effects of root-knot nematode biocontrol agent *Paecilomyces lilacinus* (*P. lilacinus*) strain PL1210 on ammonia-oxidizing microorganisms and fungal community composition of tomato rhizosphere. The exchangeable NH₄⁺-N and NO₃⁻-N contents were lower in inoculated soils than in the control during 60 days of incubation. Real-time quantitative polymerase chain reaction (qPCR) detected stable colonization of *P. lilacinus* in the tomato rhizosphere and significant inhibition of ammoniaoxidizing bacteria (AOB) and archaea (AOA), which could be responsible for the decrease of NO₃⁻-N content in soil. PCR-denaturing gradient gel electrophoresis (DGGE) analysis demonstrated no significant difference in soil fungal community composition associated with the application of *P. lilacinus* as shown by Shannon–Wiener diversity index (H') and Margalef index (D). Cluster analysis showed that the composition

of rhizosphere fungal community was more significantly influenced by time-related differences than by the inoculation of biocontrol agents.

IIM 7.1.2 Water

EU-Dossier: Doc M-IIB, Point 7.1.2

P. lilacinus is a common soil saprophyte with ubiquitous distribution worldwide. Spores of this species may also be found and persist in natural waters, but will be subject to sedimentation and do not find conditions favourable for germination and growth in this compartment. This also is supported by a test on brine shrimp, exposed to up to 8.5×10^8 CFU in 200 mL of water (██████████ 2000, M-490114-01-1): Brine shrimp were not infected, although spores were recovered from dead shrimp due to ingestion before time of death (see Doc.M, Section 6, Point 8.8, EU-Dossier Doc.M, Section 6, Point 8.4). Further, spores did not germinate in seawater alone.

Exposure to natural UV-light will to some extent restrict germination and survival of *P. lilacinus* strain 251 when spread into other environmental compartments based on the results for strain 251 tested for its UV sensitivity among other *P. lilacinus* strains (Guna-Gera et al., 2000, M-489366-01-1).

New data 2015

Statement on potential interferences with the analytical systems for the control of the quality of drinking water provided for directive 98/83/EC

According to Council Directive 98/83/EC for drinking water only *E. coli*, enterococci and *Clostridium perfringens* are monitored in drinking water. For these bacteria, either highly specific media will be used on which other species do not grow or a highly specific reaction is catalysed by the indicator species allowing a clear identification. These methods were designed to unambiguously determine these pathogens. Methods used were validated for a number of bacteria and fungi. There is no reason to assume that *P. lilacinus* would be able to grow on these media or will not catalyse the specific indicator reaction and thus interfere with the detection method.

Similarly, non-pathogenic bacteria and fungi naturally occur in drinking water systems and it has never been reported that the presence of these bacteria interferes with the quality control systems. In conclusion, any kind of interference with the analytical systems for drinking water control can be excluded.

A literature search performed in 2015 revealed three publications considered supporting for the dossier. ██████████ et al. (2010, M-534847-01-1) investigated sewage sludge. *P. lilacinus* was found among the predominating species in the sludge samples (10% of the total number of strains). These results indicate that *P. lilacinus* is a commonly occurring fungus in water systems. However, as mentioned above, spores can persist and be translocated in water compartments, but this species will rarely germinate in water. Water potential, light and pH were shown to affect the growth and sporulation of fungi (██████████, ██████████, 2009; M-534834-01). However these factors may differently affect different strains of the species. For example, for *P. lilacinus* strain IPC-P optimal water potential for growth was -7.3 MPa and for sporulation -0.3 MPa, while another strain, M-14, grew and sporulated the most efficient at -0.3 MPa. Light inhibited growth of IPC-P, but enhanced the sporulation of M-14.

Maximum growth and sporulation occurred on acid media. ██████████ et al. (2013, M-534747-01-1) investigated fungal biofilms forms on water taps in Germany. *P. lilacinus* was detected in only one out of 13 studies biofilms, and in extremely low amounts (0.23% of total fungi present). This suggests that tap water is not an optimal habitat for growth of *P. lilacinus* and thus the risk can be considered negligible.

In addition, it has to be considered that the intended application methods for BioAct WG secure a minimal drift of spores into natural surface waters (see Doc. D1).

Cited references (abstracts):

Report: KIIM 7.1.2/03 – [REDACTED], K., [REDACTED], G., [REDACTED], [REDACTED], A., [REDACTED], K., [REDACTED], S. (2010), Keratinolytic and Non-Keratinolytic Fungi in Sewage Sludge. published report

Polish J. Environ Stud 19: 635-642

Abstract: Sewage sludge is being used for reclamation of devastated areas and for fertilization of arable soils. However, sludges contain many harmful components including pathogenic organisms. Many keratinolytic and associated non-keratinolytic fungi are opportunistic pathogens. Our knowledge on fungal occurrence in sludges and sludge-amended soils and on the health risk posed by the fungi is still not sufficient. The present work was part of extensive studies on actidione-resistant fungal pathogens in sludges and sludge-amended soils. Sludges from the Siemianowice-Centrum wastewater treatment plant, Upper Silesia, Poland were examined. Results obtained by means of three methods, i.e. the dilution pour plating method, hair baiting method and most probable number method were compared. The MPN method combines the dilution and hair baiting methods. The dilution pour plating method was found not to be highly informative as to the occurrence of keratinolytic fungi in sludges, while using the method more information was obtained on non-keratinolytic fungi in the sludge environment. Subsequently the hair baiting method provided the data on fungal growth in the hair spread over the sludge blanket. This qualitative method has often been used for semiquantitative purposes but does not allow for determining fungal quantities. Such quantities were obtained using the MPN method. The method complemented the results obtained using two other methods. The hair baiting and MPN methods use hair and natural media (sterile sludge, sand, and clay) for examination of sludge fungi. The selectiveness of the MPN method was even higher than that of the hair baiting method. Ecological and epidemiological significance of the MPN results was discussed.

Report: KIIM 7.1.2/04 – [REDACTED], L., [REDACTED], X-Z., [REDACTED], M-H., [REDACTED], S-D., [REDACTED], J-L. (2009), Use of a novel two-stage cultivation method to determine the effects of environmental factors on the growth and sporulation of several biocontrol fungi. published report

Mycoscience 50: 317-321

Abstract: To supply essential information for improving mass production and biocontrol efficacy, two-stage cultivation on agar plates was used to evaluate the environmental conditions affecting mycelial growth and sporulation of seven biocontrol fungi. Maximum growth and sporulation occurred on acid media for *Paecilomyces (Po.) lilacinus* IPC-P, *Pochonia (Po.) chlamydosporia* HSY-12-9, and *Lecanicillium lecanii* CA-1-G, and on alkaline media for *Metarhizium anisopliae* isolates. All fungi preferred a certain water potential and temperature for sporulation. Light greatly inhibited the growth of *P. lilacinus* IPC-P, *M. anisopliae* SQZ-1-21, and *L. lecanii* CA-1-G but enhanced the sporulation of *P. lilacinus* M-14, *P. chlamydosporia* HSY-12-14, and *L. lecanii* CA-1-G.

Report: KIIM 7.1.2/05 – [REDACTED], G., [REDACTED], I., [REDACTED], C.K., [REDACTED], G.S., [REDACTED], G. (2015), Analysis of Black Fungal Biofilms Occurring at Domestic Water Taps (I): Compositional Analysis Using Tag-Encoded FLX Amplicon Pyrosequencing. published report

Mycopathologia 175: 387-397

Abstract: Mass growth of dark fungal biofilms on water taps and associated habitats was observed in various German drinking water distribution systems recently. Customers of affected drinking water systems are anxious about potential and unknown health risks. These environments are known to harbor a fungal flora also comprising a variety of fungal opportunists that are well known to cause superficial mycoses in humans (*Exophiala equina*, *Exophiala lecanii-corni*) but are not known to establish dark biofilms so far. To gain profound insight on composition of respective biofilms, a metagenomic approach using Tag-Encoded FLX Amplicon Pyrosequencing (TEFAP) of the ribosomal internal transcribed spacer 2 region in comparison with a classical cultivation approach using Sabouraud agar with chloramphenicol and erythritol-chloramphenicol-agar was performed. *E. lecanii-corni* was found to be the major component in 10 of 13 biofilms analyzed independently of the method used. *Alternaria* sp., *E. equina*, *Fusarium* spp. and *Ochroconis* spp. were also relatively abundant. As expected, TEFAP usually revealed a higher diversity than the cultivation approaches. For example, opportunistic species like *Candida albicans* or *Exophiala dermatitidis* were detected in very low amounts. In conclusion, TEFAP turned out to be a

promising and powerful tool for the semi-quantitative analysis of fungal biofilms. Referring to relevant literature, potential biological hazards caused by fungi of the dark biofilms can be regarded as low.

IIM 7.1.3 Air

EU-Dossier: Doc M-IIB, Point 7.1.3

Dispersal of spores via aerosols is not anticipated due to the nature of this preparation. Further information on the persistence in air is not required, since the toxicological studies and the temperature growth profile of this strain prove that it is not able to infect humans and imposes no risk for workers, operators or bystanders via the inhalation route or any other route.

IIM 7.2 Other/special studies

No further studies are considered to be necessary.

*This document is copyright protected.
Any distribution, reproduction or publication requires
the consent of Bayer AG (or its respective affiliate).
Any use of the document or its content for regulatory or
any other commercial purpose is prohibited and constitutes
a violation of the underlying license agreement.*

References

Annex point / reference number	Author(s)	Year	Title Source (where different from company) Company name, Report No., Date, GLP/GEP status (where relevant), published or not	Data protect. claimed	Owner
KIIM 7.1 /01	[REDACTED], J.; [REDACTED], J.; [REDACTED], T.; [REDACTED], S.; [REDACTED], A. M.; [REDACTED], N. L.; [REDACTED], R. A.	2011	Purpureocillium, a new genus for the medically important Paecilomyces lilacinus Year:2011, Report No.: M-534512-01-1, Edition Number: M-534512-01-1 Date: 2011-12-31 GLP/GEP: no, published ...also filed: KIIM 1.3.1 /01 ...also filed: KIIM 1.3.3 /05 ...also filed: KIIM 2.7.1 /18 ...also filed: KIIM 5.2.4 /03 ...also filed: KIIM 6.1 /02	No	
KIIM 7.1 /02	[REDACTED], A.	2015	Literature review on Purpureocillium lilacinum strain 251: Fate and behaviour in the environment G&B Consulting Asociencias S.L.U. Valencia, Spain. Bayer CropScience, Report No.: M-542801-01-1, Edition Number: M-542801-01-1 Date: 2015-12-17 GLP/GEP: n.a., unpublished	Yes	Bayer CropScience
KIIM 7.1.1 /01	[REDACTED], L. P.-H.	1985	Persistence des conidies spores des hyphomycetes entomopathogenes bouveria (assiana) vuill., metarhizium anisopliae metsch.) sor., nomurea rileyi (f.) samson et paecilomyces fumoso-roseus wize dans le sol, en conditions controlees Journal: agronomie, Volume: 1, Issue: 1, Pages:73-80, Year:1985, Report No.: M-489363-01-1, Edition Number: M-489363-01-1 GLP/GEP: n.a., published ...also filed: KIIM 2.8 /10	No	
KIIM 7.1 /02	[REDACTED], R. D.; [REDACTED], J.	1990	Relationship between inoculum density of the nematophagous fungus Paecilomyces lilacinus and control of Meloidogyne arenaria on tomato Journal:Revue Nématol., Volume:14, Issue:4, Pages:629-634, Year:1991, Report No.: M-477635-01-1, Edition Number: M-477635-01-1 GLP/GEP: n.a., published ...also filed: KIIM 2.8 /11	No	

Annex point / reference number	Author(s)	Year	Title Source (where different from company) Company name, Report No., Date, GLP/GEP status (where relevant), published or not	Data protect. claimed	Owner
KIIM 7.1.1 /03	[REDACTED], R.K.; [REDACTED], S.N.; [REDACTED] D.S.	1998	Nematicidal efficacy of plant leaves and Paecilomyces lilacinus, alone or in combination, in controlling Meloidogyne incognita on okra and tomato Journal:Nematol. medit., Volume:27, Pages:3-8, Year:1999, Report No.: M-477445-01-1 Edition Number: M-477445-01-1 GLP/GEP: n.a., published ...also filed: KIIM 2.1 /12	No	
KIIM 7.1.1 /04	[REDACTED], Z.A.; [REDACTED] I.	1996	Biological control of plant parasitic nematodes by fungi: A review Publisher:Elsevier Science Ltd, Location:Great Britain, Journal:Bioresource Technology, Volume:5, Pages:229-239, Year:1996, Report No.: M-477546-01-1, Edition Number: M-477546-01-1 GLP/GEP: n.a., published ...also filed: KIIM 2.1 /01 ...also filed: KIIM 2.2 /01 ...also filed: KIIM 2.7 /03 ...also filed: KIIM 2.8 /13	No	
KIIM 7.1.1 /05	[REDACTED], P.	1986	Biological control of plant-parasitic nematodes Publisher:Annual Reviews Inc., Journal:Ann. Rev.Phytopathol., Volume: 4, Pages:453-489, Year:1986, Report No.: M-477553-01-1, Edition Number: M-477553-01-1 GLP/GEP: n.a., published ...also filed: KIIM 2.1 /03 ...also filed: KIIM 2.3.1 /04 ...also filed: KIIM 2.3.2 /02 ...also filed: KIIM 2.5 /04 ...also filed: KIIM 2.8 /02 ...also filed: KIIM 3.4 /04 ...also filed: KIIM 3.4.2 /04	No	

This document is copyright protected. Any distribution, reproduction or publication requires the consent of Bayer AG (or its respective affiliate). Any use of the document for regulatory or commercial purposes is prohibited and constitutes a violation of the underlying license agreement.

Annex point / reference number	Author(s)	Year	Title Source (where different from company) Company name, Report No., Date, GLP/GEP status (where relevant), published or not	Data protect. claimed	Owner
KIIM 7.1.1 /06	[REDACTED], P.; [REDACTED], R.; [REDACTED], R.; [REDACTED] [REDACTED], M.	1981	Multiple application and long-term effects of Paecilomyces lilacinus in controlling Meloidogyne incognita under field conditions Publisher:Society of Nematologists, Journal:Journal of Nematology, Annual Abstracts, Volume:13, Pages:445, Year:1981 Report No.: M-477590-01-1 Edition Number: M-477590-01-1 GLP/GEP: n.a., published ...also filed: KIIM 2.1 /08 ...also filed: KIIM 2.8 /14	No	
KIIM 7.1.1 /07	[REDACTED], C.B.; [REDACTED], R.M.D.G.	1997	Receptividade de diferentes solos a colonizaco pelo fungo paecilomyces lilacinus em reses formicoes. Volume:2 Issue:2, Pages:61-73, Year:1997, Report No.: M-489365-01-1, Edition Number: M-489365-01-1 GLP/GEP: n.a., published ...also filed: KIIM 2.1 /15	No	
KIIM 7.1.1 /08	[REDACTED], E.; [REDACTED], K. R.; [REDACTED], L. A.	1989	Survival of paecilomyces lilacinus in selected carriers and related effects on meloidogyne incognita on tomato Publisher:The Society of Nematology, Journal:Journal of Nematology, Volume:1, Issue:1, Pages:121-130, Year:1989, Report No.: M-489356-01-1, Edition Number: M-489356-01-1 GLP/GEP: n.a., published	No	
KIIM 7.1.1 /09	[REDACTED], K.H.; [REDACTED], W.; [REDACTED] [REDACTED], T.	1993	Compendium of soil fungi - Volume 1 Publisher:IHW-Verlag, Journal:Compendium of soil fungi, Volume:1, Pages:529-532, Year:1993, Report No.: M-476530-01-1, Edition Number: M-476530-01-1 GLP/GEP: n.a., published ...also filed: KIIM 1.3.1 /04 ...also filed: KIIM 2.2 /02 ...also filed: KIIM 2.5 /02 ...also filed: KIIM 2.8 /01	No	

Annex point / reference number	Author(s)	Year	Title Source (where different from company) Company name, Report No., Date, GLP/GEP status (where relevant), published or not	Data protect. claimed	Owner
KIIM 7.1.1 /10	[REDACTED] V.M.; [REDACTED] T.K.	1984	Effect of soil fungistasis on zoopathogenic fungi Publisher:Dr W. Junk Publishers, The Hague, Location:The Netherlands, Journal:Mycopathologia, Volume:85, Pages:121-122 Year:1984, Report No. M-477420-01-1 Edition Number: M-477420-01-1 GLP/GEP: n.a. published ...also filed: KIIM 2.8 /16	No	
KIIM 7.1.1 /11	[REDACTED] R.	1998	Interim report - Persistence of Paecilomyces lilacinus in garden soil Macquarie University, Sydney, Australia Bayer CropScience, Report No. M-477414-01-1 Edition Number: M-477414-01-1 Date: 1998-02-25 GLP/GEP: n.a. unpublished	Yes	Bayer CropScience
KIIM 7.1.1 /12	[REDACTED] R.	2000	Paecilomyces lilacinus as a biocontrol agent - Chapter 7 - The persistence of Paecilomyces lilacinus after application to soil Report No.: M-490110-01-1, Edition Number: M-490110-01-1 GLP/GEP: n.a. published ...also filed: KIIM 2.8 /17 ...also filed: KIIM 4.5.4 /01	No	
KIIM 7.1.1 /13	[REDACTED]	2005	Statement to the DAR reference point: Vol. 3 - 8.1, persistence and multiplication in soil Prophyta, Malchow/Poel, Germany Bayer CropScience, Report No.: M-542625-01-1, Edition Number: M-542625-01-1 Date: 2005-12-14 GLP/GEP: n.a., unpublished	Yes	Bayer CropScience
KIIM 7.1.1 /15	[REDACTED] S.; [REDACTED] R.	2006	Biological control of the root-knot nematode Meloidogyne incognita by Paecilomyces lilacinus strain 251 Journal:Biological control : theory and application in pest management (2006) , Year:2006, Report No.: M-534735-01-1, Edition Number: M-534735-01-1 Date: 2006-12-31 GLP/GEP: no, published	No	

Annex point / reference number	Author(s)	Year	Title Source (where different from company) Company name, Report No., Date, GLP/GEP status (where relevant), published or not	Data protect. claimed	Owner
KIIM 7.1.1 /16	[REDACTED], C. I.; [REDACTED], S.	2006	Effect of plant species on persistence of Paecilomyces lilacinus strain 251 in soil and on root colonization by the fungus Year:2006, Report No.: M-534361-01-1 Edition Number: M-534361-01-1 Date: 2006-12-31 GLP/GEP: no, published ...also filed/KIIM 6.1 /04	No	
KIIM 7.1.1 /17	[REDACTED], S.; [REDACTED], C.; [REDACTED], R. A.	2005	Risk assessment of biological control products. Risikoabschätzung von biologischen Pflanzenschutzmitteln: Persistenz von Paecilomyces lilacinus Stamm 251 unter Freilandbedingungen. Journal: Gesunde Pflanzen (2005), Year:2005, Report No. M-534360-01-1 Edition Number: M-534360-01-1 Date: 2005-12-31 GLP/GEP: no, published	No	
KIIM 7.1.1 /18	[REDACTED], S.; [REDACTED], C.; [REDACTED], R. E. P. B.	2005	Risk assessment of biological nematocides. Journal: Bulletin OILB/SROP (2005), Year:2005, Report No.: M-535175-01-1, Edition Number: M-535175-01-1 Date: 2005-12-31 GLP/GEP: no, published	No	
KIIM 7.1.1 /19	[REDACTED], R.; [REDACTED], E.; [REDACTED], W.	2015	Modifying soil to enhance biological control of belowground dwelling insects in citrus groves under organic agriculture in Florida. Journal: Biological Control (2015), Year:2015, Report No.: M-534528-01-1, Edition Number: M-534528-01-1 Date: 2015-09-23 GLP/GEP: no, published	No	
KIIM 7.1.1 /20	[REDACTED], S. E. R. J. M.; [REDACTED], R. A.	2006	Multitrophic interactions of Paecilomyces lilacinus strain 251 in the rhizosphere of host and non-host plants. Journal: Bulletin OILB/SROP (2006), Year:2006, Report No.: M-534359-01-1, Edition Number: M-534359-01-1 Date: 2006-12-31 GLP/GEP: no, published	No	

Annex point / reference number	Author(s)	Year	Title Source (where different from company) Company name, Report No., Date, GLP/GEP status (where relevant), published or not	Data protect. claimed	Owner
KIIM 7.1.1 /21	[REDACTED], C.; [REDACTED], A.; [REDACTED], R.; [REDACTED], S.	2008	Persistence of the nematophagous fungus Paecilomyces lilacinus strain 251 in soil under controlled conditions. Journal: Biocontrol science and technology (2008), Year: 2008, Report No.: M-534367-01-1, Edition Number: M-534367-01-1 Date: 2008-12-31 GLP/GEP: no, published	No	
KIIM 7.1.1 /22	[REDACTED], S. E. E. Y.; [REDACTED], M.; [REDACTED], C.; C.; [REDACTED], B.	2009	Understanding multitrophic interactions to facilitate successful biocontrol of plant parasitic nematodes with Paecilomyces lilacinus strain 251. Journal: IOBC/WPRS Bulletin (2009) Year: 2009 Report No.: M-534891-01-1, Edition Number: M-534891-01-1 Date: 2009-12-31 GLP/GEP: no, published	No	
KIIM 7.1.1 /23	[REDACTED], M. J. A.; [REDACTED], P. P.	2005	Management of carnation and gerbera to control the root-knot nematode, Meloidogyne incognita in commercial polyhouses. Year: 2005 Report No.: M-534724-01-1, Edition Number: M-534724-01-1 Date: 2005-12-31 GLP/GEP: no, published	No	
KIIM 7.1.1 /24	[REDACTED], D. S.; [REDACTED], M. S.; A.; [REDACTED] [REDACTED], R. M.; [REDACTED], J.; [REDACTED] K.	2012	BIO-MANAGEMENT OF MELOIDOGYNE INCOGNITA AND ERWINIA CAROTOVORA IN CARROT (DAUCUS CAROTA L.) USING PSEUDOMONAS PUTIDA AND PAECILOMYCES LILACINUS . Year: 2012, Report No.: M-534851-01-1, Edition Number: M-534851-01-1 Date: 2012-12-31 GLP/GEP: no, published	No	

Annex point / reference number	Author(s)	Year	Title Source (where different from company) Company name, Report No., Date, GLP/GEP status (where relevant), published or not	Data protect. claimed	Owner
KIIM 7.1.1 /25	Z.; Y.; W.; Y.	2015	Root colonization and effect of biocontrol fungus <i>Paecilomyces lilacinus</i> on composition of ammonia-oxidizing bacteria, ammonia-oxidizing archaea and fungal populations of tomato rhizosphere Year:2015, Report No.: M-534211-01-1, Edition Number: M-534211-01-1 Date: 2015-09-25 GLP/GEP: no, published	No	
KIIM 7.1.1 /14	T.S.; R.J.; MR.; D.A.; D.C.; K.L.; K.M.H.	2000	Phenotypic and genetic characterization of <i>paecilomyces lilacinus</i> strains with biocontrol activity against root-knot nematode Publisher: NRC Canada, Journal: Can. J. Microbiol., Volume: 46, Pages: 77-83, Year: 2000, Report No.: M-489366-01-1, Edition Number: M-489366-01-1 GLP/GEP: n.a., published ...also filed: KIIM 1.3.1 /03 ...also filed: KIIM 1.3.1 /03 ...also filed: KIIM 2.1 /01 ...also filed: KIIM 2.1 /07 ...also filed: KIIM 4.1 /02 ...also filed: KIIM 4.3.1 /02 ...also filed: KIIM 7.1.2 /02	No	
KIIM 7.1.2 /01	R. J.	2001	<i>Paecilomyces lilacinus</i> as a biocontrol agent - Chapter 6 - The effect of <i>Paecilomyces lilacinus</i> on non-target invertebrates Report No.: M-490114-01-1, Edition Number: M-490114-01-1 GLP/GEP: n.a., published ...also filed: KIIM 8.3 /02 ...also filed: KIIM 8.8 /03 ...also filed: KIIM 8.9.2 /02	No	

This document is copyright protected. Any distribution, reproduction or publication requires the consent of Bayer AG (or its respective affiliate). Any use of the document for regulatory or commercial purposes is prohibited and constitutes a violation of the underlying license agreement.

Annex point / reference number	Author(s)	Year	Title Source (where different from company) Company name, Report No., Date, GLP/GEP status (where relevant), published or not	Data protect. claimed	Owner
KIIM 7.1.2 /02	[REDACTED], T.S.; [REDACTED] R.J.; [REDACTED] M.R.; [REDACTED] D.A.; [REDACTED] D.C.; [REDACTED] K.L.; [REDACTED] K.M.H.	2000	Phenotypic and genetic characterization of paecilomyces lilacinus strains with biocontrol activity against root-knot nematodes Publisher:NRC Canada, Journal:Can. J. Microbiol. Volume:46, Pages:775-782 Year:2000, Report No.:M-489366-01-1 Edition Number: M-489366-01-1 GLP/GEP: n.a. published ...also filed: KIIM 1.1 /03 ...also filed: KIIM 1.3 /03 ...also filed: KIIM 2.1 /01 ...also filed: KIIM 2.8 /01 ...also filed: KIIM 4.1 /02 ...also filed: KIIM 4.1 /02 ...also filed: KIIM 4.1 /02	No	
KIIM 7.1.2 /03	[REDACTED] K. J. A.; [REDACTED] G.; [REDACTED] A.; [REDACTED] K.; [REDACTED] S.	2010	Keratinolytic and Non-Keratinolytic Fungi in Sewage Sludge Year:2010, Report No.: M-534847-01-1, Edition Number: M-534847-01-1 Date:2010-12-31 GLP/GEP: no, published	No	
KIIM 7.1.2 /04	[REDACTED] I.; [REDACTED] A.; [REDACTED] M.; [REDACTED] S.; [REDACTED] J.	2009	Use of a novel two-stage cultivation method to determine the effects of environmental factors on the growth and sporulation of several biocontrol fungi. Year:2009, Report No.: M-534834-01-1, Edition Number: M-534834-01-1 Date: 2009-12-31 GLP/GEP: no, published	No	
KIIM 7.1.2 /05	[REDACTED] G.; [REDACTED] I.; [REDACTED] C.; [REDACTED] S.; [REDACTED] G.	2013	Analysis of Black Fungal Biofilms Occurring at Domestic Water Taps (I): Compositional Analysis Using Tag-Encoded FLX Amplicon Pyrosequencing Year:2013, Report No.: M-534747-01-1, Edition Number: M-534747-01-1 Date: 2013-12-31 GLP/GEP: no, published	No	